This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

HN 25V5 /

WHO'S·WHO AMONG·THE WILD·FLOWERS

W. I. BEECROFT

Google

KD 5647-6

HARVARD COLLEGE
LIBRARY

THE BEQUEST OF

EVERT JANSEN WENDELL
(CLASS OF 1882)

OF NEW YORK

1918

Digitized by Google

Digitized by Google

Digitized by Google

WHO'S WHO
AMONG THE
WILD FLOWERS

Digitized by Google

Digitized by Google

WHO'S WHO AMONG THE WILD FLOWERS

COMPILED AND ILLUSTRATED

BY

W. I. BEECROFT

WITH AN INTRODUCTION

BY

FRANCES DUNCAN

NEW YORK
MOFFAT, YARD AND COMPANY
1911

Digitized by Google



KD 3845 G

HARVARD COLLEGE LIBRARY
FROM
THE BEQUEST OF
EVERT JANSEN WENDELL
1918

Copyright 1910, by
MOFFAT, YARD AND COMPANY
NEW YORK

All Rights Reserved

Published January, 1910
Reprinted March, 1911

Digitized by Google

INTRODUCTION

There may be perverse and misguided folk, who, like myself, have an unaccountable distaste for looking at Mother Nature through the spectacles of Another. One is humbly aware that the "spectacles" offered by the various writers are admirable; that they clarify and enlarge the vision, place the object in an excellent light and show our much-inspected Mother as wearing the loveliest of complexions, and yet—one likes to use his own eyesight—his own spectacles. To such, this little book will prove a peculiar blessing. It goes no farther than to tell "Who's Who" among the wild flowers, merely giving an introduction and allowing one the privilege of making friends with them after the devices of his own heart.

Mr. Beecroft believes that a portrait if an excellent likeness and true in all its details will enable the most unskill'd, unpractis'd and unbotanical person to recognize his new acquaintance more speedily than much writing of many paragraphs. The drawings are supplemented by the briefest of notes on the personal appearance—only enough to identify, and in the blank spaces "remarks" can be made by the reader who is allowed the chance of making a flower book after the devices of his own

Digitized by Google

heart; thus an idler will make it the record of a summer's wanderings and write in bits of verse and prose that fit the subjects; the exact and botanical soul can add descriptions, Bertillon measurements, variations of nomenclature, drawings of parts to heart's content: children will enjoy mounting the flowers they find opposite their pictures. While to the botanist it will be a most convenient field-book, for at the end of the season his various jottings will be found already classified and indexed.

In arrangement, the excellent method inaugurated by Mrs. Parsons, now customary in most non-technical flower-books, has been followed here, that of grouping the flowers according to color. Within the color-divisions the plants are arranged according to their time of appearance.

The nomenclature is that of the latest edition of Gray's "Manual." The pronunciation is indicated, since it sacrificed no space and may be a convenience to children and those of us whose instincts about the pronunciation of a botanical name are not always infallible.

The common names of plants are legion, of these, the best known and most generally used are in large type, the less important ones in small type and they are given lest some reader look for a flower under the alias by which he knows it, and missing this should think his friend omitted in this "Who's Who."

Digitized by Google

INTRODUCTION

It may be well to remind flower-loving folk that some of our plants are in sore need of a little intelligent affection. Unless speedily befriended, the exquisite Trailing Arbutus will quite vanish from our hillsides, leaving behind only a memory of its loveliness. The cutting of the blossoms would be harmless enough, but it is the ripping up of the creeping, slow-growing root-fibres for convenient picking which makes the extinction of this darling New-Englander so certain. The bunches for sale in the city streets represent the death of millions of little unborn Mayflowers. The next plant to go will be the Mountain Laurel. Other flowers whose existence is in peril are the Fringed Gentian and Hepatica, Orchids, Maiden Hair Fern, Sabatia, Ground Pine or Club Moss and the Hollies.

The nature love which manifests itself in a kind of pot-hunter's enthusiasm or sees in each rare flower only a kind of botanical scalp to be added to his belt, is a poor sort of affection, and though perhaps one cannot sing about that Millennium of Flowers when folk will be content to love the wood rose and leave it on its stalk, we may live to see it cut instead of torn from its stem and the last Fringed Gentian respected and left to perpetuate its lovely kind.

FRANCES DUNCAN.

BROOKLYN, NEW YORK, January, 1910.

Digitized by Google

Digitized by Google

CONTENTS.

INTRODUCTION v

I

WHITE AND GREENISH

PAGE

BLOODROOT 2
DUTCHMAN'S BREECHES 4
SQUIRREL CORN 4
TOOTHWORT 6
CUT-LEAVED TOOTHWORT 6
PARTRIDGE BERRY 6
FOAM-FLOWER 10
MITREWORT 10
NAKED MITREWORT 10
EARLY SPRING EVERLASTING 12
EARLY SAXIFRAGE 12
RED BANEBERRY 14
WHITE BANEBERRY 14
SOLOMON'S SEAL 16
GREAT SOLOMON'S SEAL 16
RUE ANEMONE 18
WOOD ANEMONE 18
WILD SARSAPARILLA 20
DWARF GINSENG 20
ARROW ARUM 24
WATER ARUM 26
MAIANTHEMUM 26
PAINTED TRILLIUM 28
NODDING TRILLIUM 28
LARGE-FLOWERED TRILLIUM 28

Digitized by Google

PAGE

Star Flower 30
Goldthread 30
False Solomon's Seal 32
Star-flowered Solomon's Seal 32
Poison Sumach 34
Poison Ivy 34
Bristly Sarsaparilla 36
Shin Leaf 38
Round-leaved Pyrola 38
One-flowered Pyrola 38
Red-berried Elder 40
Common Elder 40
Sweet Pepperbush 40
Rein Orchis 42
Thimbleweed 44
Field Chickweed 44
Indian Pipe 46
Wild Leek 46
Bladder Campion 50
White Campion 50
Musk Mallow 52
Common Mallow 52
Chamomile 54
Bunchberry 54
Daisy Fleabane (2) 56
Yarrow 58
Virgin's Bower 60
Ragged Fringed Orchis 62
Rattlesnake Plantain 62
Greater Green Orchis 64
Hooker's Orchis 64
Round-leaved Sundew 66
Long-leaved Sundew 66
Slender Ladies' Tresses 66
Thoroughwort 68

Digitized by Google

PAGE

White Snakeroot 68
Turtlehead 70
Arrow-head 72
Rough Bedstraw 72
Wild Balsam-apple 76
Tall Meadow Rue 78
Grass of Parnassus 80
Water Plantain 82
Common Dodder 84
Wood Sorrel 84
Pearly Everlasting 86
White Melilot 86
Wild Carrot 88
Buttonbush 90
Pokeweed 92
Flowering Moss 94
Mayflower 94
White Adder's Tongue 96
Twinleaf 96
May Apple 97
Sweet White Violet 97
Lance-leaved Violet 98
Canada Violet 98
Water Cress 99
Creeping Snowberry 99
New Jersey Tea 100
Black Snakeroot 100
Clammy Azalea 100
Showy Lady's Slipper 101
Lizard's Tail 101
Poke Milkweed 102
Water Hemlock 102
Water Parsnip 102
Dalibarda 103
Water Lily 103

Digitized by Google

PAGE

White Fringed Orchis 104
Prairie White Fringed Orchis 104
Wintergreen 104
Culver's Root 105
Jimson Weed 105

II

YELLOW AND ORANGE

Yellow Adder's Tongue 110
Marsh Marigold 112
Star Grass 114
Cinquefoil 114
Golden Alexanders 116
Wild Parsnip 116
Clintonia 118
Indian Cucumber-root 120
Oakesia 122
Large-flowered Bellwort 122
Bellwort 122
Yellow Lady's Slipper 126
Small Yellow Lady's Slipper 126
Golden Ragwort 128
Carrion-flower 130
Yellow Pond Lily 132
Wild Yellow Lily 134
Turk's-cap Lily 134
Fringed Loosestrife 136
Agrimony 138
Celandine 138
Loosestrife 140
Four-leaved Loosestrife 140
Cone-flower 142
Tall Cone-flower 142
Common Mullein 144

Digitized by Google

PAGE

Moth Mullein 144
Wild Lettuce 146
Yellow, or Hop Clover 148
Lady's Sorrel 148
Black Mustard 150
Yellow Rocket 150
Common Evening Primrose 154
Sundrops 154
Spotted Touch-me-not 156
Pale Touch-me-not 156
Common St. John's Wort 158
Butter-and-Eggs 160
Yellow Fringed Orchis 162
Downy False Foxglove 162
Smooth False Foxglove 162
Elecampane 164
Partridge Pea 166
Wild Senna 166
Fall Dandelion 166
Tansy 170
Orange Hawkweed 172
Canada Hawkweed 172
Hawkweed 174
Hairy Hawkweed 174
Rattlesnake-weed 174
Thin-leaved Sunflower 176
Larger Bur-marigold 176
Golden Corydalis 179
Round-leaved Violet 179
Downy Yellow Violet 180
Golden Club 180
Indian Poke 181
Yellow Melilot 181
Frostweed 182
Butterfly Weed 183

Digitized by Google

PAGE

Ginseng 183
Horse Balm 184
Sneezeweed 184
Fern-leaved False Foxglove 185
Golden Aster 185
Golden Rod 185

III

RED, PINK, CRIMSON AND MAGENTA

Spring Beauty 188
Shooting Star 190
Arethusa 190
Wake Robin 192
Sessile-flowered Wake Robin 192
Wild Columbine 194
Wild Pink 196
Maiden Pink 196
Deptford Pink 196
Wild Cranesbill 198
Herb Robert 198
Pitcher-plant 202
Stemless Lady's Slipper 204
Showy Orchis 206
Wild Azalea 208
Fringed Milkwort 210
Field Milkwort 210
Mountain Laurel 212
Twisted Stalk 214
Clasping-leaved Twisted-stalk 214
Wood Betony 216
Bouncing Bet 216
Pale Corydalis 218
Sheep Laurel 220
Pale Laurel 220

Digitized by Google

PAGE

Grass Pink 222
Rose Pogonia 222
Wild Orange-red Lily 224
Spreading Dogbane 226
Twin-flower 226
Meadow-sweet 228
Hardhack 228
Purple Flowering Raspberry 230
Purple Fringed Orchis 234
Smaller Purple Fringed Orchis 234
Prince's Pine 234
Spotted Wintergreen 234
Common Milkweed 236
Purple Milkweed 236
Hedge Bindweed 238
Field Bindweed 238
Large Marsh Pink 240
Bitter Bloom 240
Sea Pink 240
Slender Marsh Pink 240
Swamp Milkweed 242
Four-leaved Milkweed 242
Great Willow-herb 244
Cardinal Flower 244
Meadow Beauty 246
Maryland Meadow Beauty 246
Oswego Tea 246
Pink Knotweed 248
Lady's Thumb 248
Water Persicaria 248
Joe-Pye Weed 250
Rabbit's-foot Clover 252
Live-for-ever 254
Ground Pink 258
Downy Phlox 258

Digitized by Google

PAGE

Violet Wood Sorrel 259
Hairy Beard-tongue 259
Painted Cup 260
Queen of the Prairie 261
Common Pasture Rose 262
Swamp Rose 262
Early Rose 262
Sweet-brier; Eglantine 263
Wild Bergamot 263
Purple Cone-flower 264
Cross-leaved Milkwort 264
Corn Cockle 265
Swamp Rose Mallow 265
Purple Gerardia 266

IV

BLUE AND PURPLE

Liverleaf 270
Sharp-lobed Liverleaf 270
Bird-foot Violet 272
Robin's Plantain 274
Wild Lupine 274
Bluets 276
Gill-over-the-Ground 276
Larger Blue Flag 278
Slender Blue Flag 278
Cow Vetch 280
American Vetch 280
Bittersweet; Nightshade 282
Self-heal 284
Blue-eyed Grass 284
Bellflower 286
Harebell 286
Pale Spiked Lobelia 290

Digitized by Google

PAGE

Motherwort 290
Pickerel-weed 292
Monkey Flower 292
Mad-dog Skullcap 294
Skullcap 294
Blue Vervain 296
Wild Mint 298
Spearmint 298
Peppermint 298
Ironweed 300
Chicory 302
Blazing Star 302
New England Aster 304
Fringed Gentian 306
Closed Gentian 306
Virginian Cowslip 310
Early Blue Violet 310
Arrow-leaved Violet 311
Dog Violet 311
Common Violet 312
Wild Blue Phlox 312
Forget-me-not 313
Smaller Forget-me-not 313
Spiderwort 313
Beach Pea 314
Common Speedwell 314
American Brooklime 315
Virginia Day-flower 315
Blue Toadflax 316
Viper's Bugloss 316
Tall Wild Larkspur 317
Dwarf Larkspur 317
Purple Orchis 317
Sea Lavender 318

Digitized by Google

V

MISCELLANEOUS

	PAGE
Skunk Cabbage	322
Wild Ginger	324
Jack-in-the-Pulpit	326
Branching Bur-reed	328
Sweet Flag	328
Pinesap	330
Beechdrops	330
Large Coral Root	332
Rattlesnake-root	332
Groundnut	334
Early Meadow Rue	338
Blue Cohosh	338
Common Alum Root	339
Large Twayblade	340
Cut-tail Flag	341
Narrow-leaved Cat-tail Flag	341

Digitized by Google

LIST OF ILLUSTRATIONS

PAGE

Bloodroot 3
Dutchman's Breeches 5
Squirrel Corn 5
Toothwort 7
Partridge Berry 7
Foam-flower 11
Mitrewort 11
Early Everlasting 13
Early Saxifrage 13
Red Baneberry 15
Solomon's Seal 17
Rue Anemone 19
Wood Anemone 19
Wild Sarsaparilla 21
Dwarf Ginseng 21
Arrow Arum 25
Water Arum 27
Maianthemum 27
Painted Trillium 29
Star Flower 31
Goldthread 31
False Solomon's Seal 33
Poison Sumach 35
Poison Ivy 35
Bristly Sarsaparilla 37
Shin Leaf 39
Red-berried Elder 41
Sweet Pepperbush 41

Digitized by Google

PAGE

Rein Orchis (2) 43
Thimbleweed 45
Field Chickweed 45
Indian Pipe 47
Wild Leek 47
Bladder Campion 51
White Campion 51
Musk Mallow 53
Common Mallow 53
Chamomile 55
Bunch Berry 55
Daisy Fleabane 57
Yarrow 59
Virgin's Bower 61
Ragged Fringed Orchis 63
Rattlesnake Plantain 63
Greater Green Orchis 65
Round-leaved Sundew 67
Ladies' Tresses 67
Thoroughwort 69
Turtlehead 71
Arrow-head 73
Rough Bedstraw 73
Wild Balsam-apple 77
Tall Meadow Rue 79
Grass of Parnassus 81
Water Plantain 83
Dodder 85
Wood Sorrel 85
Pearly Everlasting 87
White Melilot 87
Wild Carrot 89
Buttonbush 91
Pokeweed 93
Yellow Adder's Tongue 111

Digitized by Google

PAGE

Marsh Marigold 113
Star Grass 115
Cinquefoil 115
Golden Alexanders 117
Wild Parsnip 117
Clintonia 119
Indian Cucumber-root 121
Oakesia 123
Large-flowered Bellwort 123
Bellwort 123
Yellow Lady's Slipper 127
Golden Ragwort 129
Carrion-flower 131
Yellow Pond Lily 133
Wild Yellow Lily 135
Fringed Loosestrife 137
Agrimony 139
Celandine 139
Loosestrife 141
Four-leaved Loosestrife 141
Cone-flower 143
Common Mullein 145
Moth Mullein 145
Wild Lettuce 147
Yellow Clover 149
Lady's Sorrel 149
Black Mustard 151
Common Evening Primrose 153
Sundrops 153
Spotted Touch-me-not 157
Common St. John's Wort 159
Butter-and-Eggs 161
Yellow Fringed Orchis 163
Downy False Foxglove 163
Elecampane 165

Digitized by Google

PAGE

Partridge Pea 167
Fall Dandelion 167
Tansy 171
Orange Hawkweed 173
Hawkweed 175
Rattlesnake-weed 175
Thin-leaved Sunflower 177
Larger Bur-marigold 177
Spring Beauty 189
Shooting Star 191
Arethusa 191
Wake Robin 193
Wild Columbine 195
Wild Pink 197
Maiden Pink 197
Wild Cranesbill 199
Herb Robert 199
Pitcher-plant 203
Stemless Lady's Slipper 205
Showy Orchis 207
Wild Azalea 209
Fringed Milkwort 211
Field Milkwort 211
Mountain Laurel 213
Twisted-stalk 215
Wood Betony 217
Bouncing Bet 217
Pale Corydalis 219
Sheep Laurel 221
Grass Pink 223
Rose Pogonia 223
Wild Orange-red Lily 225
Spreading Dogbane 227
Twin-flower 227
Meadow-sweet 229

Digitized by Google

PAGE

HARDHACK 229
PURPLE FLOWERING RASPBERRY 231
PURPLE FRINGED ORCHIS 235
PRINCE'S PINE 235
COMMON MILKWEED 237
HEDGE BINDWEED 239
LARGE MARSH PINK 241
SEA PINK 241
SWAMP MILKWEED 243
GREAT WILLOW-HERB 245
CARDINAL FLOWER 245
MEADOW BEAUTY 247
OSWEGO TEA 247
PINK KNOTWEED 249
WATER PERSICARIA 249
JOE-PYE WEED 251
RABBIT'S-FOOT CLOVER 253
LIVE-FOR-EVER 255
LIVERLEAF 271
BIRD-FOOT VIOLET 273
ROBIN'S PLANTAIN 275
WILD LUPINE 275
BLUETS 277
GILL-OVER-THE-GROUND 277
LARGER BLUE FLAG 279
COW VETCH 281
BITTERSWEET 283
SELF-HEAL 285
BLUE-EYED GRASS 285
BELLFLOWER 287
HAREBELL 287
PALE SPIKED LOBELIA 291
MOTHERWORT 291
PICKEREL-WEED 293
MONKEY FLOWER 293

Digitized by Google

PAGE

MAD-DOG SKULLCAP 295
SKULLCAP 295
BLUE VERVAIN 297
WILD MINT 299
SPEARMINT 299
PEPPERMINT 299
IRONWEED 301
CHICORY 303
BLAZING STAR 303
NEW ENGLAND ASTER 305
FRINGED GENTIAN 307
CLOSED GENTIAN 307
SKUNK CABBAGE 323
WILD GINGER 325
JACK-IN-THE-PULPIT 327
BRANCHING BUR-REED 329
SWEET FLAG 329
PINESAP 331
BEECHDROPS 331
CORAL ROOT 333
RATTLESNAKE-ROOT 333
GROUNDNUT 335

Digitized by Google

I

WHITE AND GREENISH

Digitized by Google

BLOODROOT: Indian Paint: Red Puccoon.

Sanguinària canadénsis.

Poppy Family Apr.—May.

Found in open woods and borders of woodlands. Nova Scotia to Florida, west to Nebraska.

FLOWERS—Pure white, 8–12 divisions, solitary on stalk 6-12 inches high.

LEAVES— Rounded, deeply lobed, springing from thick rootstock.

Juice of root orange colored.

(*See Frontispiece.*)

NOTES.

Digitized by Google



Bloodroot.

Digitized by Google

1. DUTCHMAN'S BREECHES: Ear-drops: White Hearts: Soldiers' Cap.
Dicéntra Cucullària.
Fumitory Family. Apr.—May.
Found in rich, rocky woods. Nova Scotia to the Carolinas, common westward.
FLOWERS—White, tipped with pale yellow, strung on slender stalk 5–10 inches high.
LEAVES— Finely cut, from a bulbous root.

2. SQUIRREL CORN: *Dicéntra canadénsis.*
Fumitory Family. Apr.—May.
Found in rich woods. Nova Scotia to Virginia, west to Mississippi.
FLOWERS—White, tinged with magenta, fragrant; 4–8 on a stalk 6–12 inches high.
LEAVES— Finely cut, from the base. Roots bear small, yellow tubers.

NOTES.

Digitized by Google



1. Dutchman's Breeches. **2. Squirrel Corn.**

Digitized by Google

1. TOOTHWORT; CRINKLE-ROOT:
Dentària diphýlla.

Mustard Family. Apr.—May.

Found in rich woods. Nova Scotia, Maine to Minnesota, south to the Carolinas, west to the Mississippi.

FLOWERS—White, 4 divisions, borne in a loose cluster terminating a stalk 8–13 inches high.

LEAVES— 2 opposite, compound leaves borne on flower stalk, leaves at base larger. Rootstalk long, crinkled, possessing pungent flavor.

2. CUT-LEAVED TOOTHWORT: Pepper-root.
D. laciniàta.

Plant similar to the above.

FLOWERS—White or pinkish.

LEAVES— Stem leaves in whorls of 3, much divided, toothed. Roots tuberous, edible.

3. PARTRIDGE BERRY: Partridge Vine: Twin-berry: Mitchella-vine: Squawberry.
Mitchélla rèpens.

Madder Family. May—June.

Found in woods. Nova Scotia to Gulf of Mexico, westward to Minnesota and Texas.

FLOWERS—White, fragrant, in pairs at the tips of the sprays. Berries red, edible.

LEAVES— Rounded, evergreen, in pairs on a trailing plant. Stems, 6–12 inches long.

Digitized by Google

1. Toothwort. **2. Partridge Berry.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

1. FOAM-FLOWER; FALSE MITREWORT: Coolwort: Nancy-over-the-ground.
Tiarélla cordifòlia.

Saxifrage Family. Apr.—June.

Found in rich woods. Nova Scotia to Georgia, westward to Minnesota and Indiana.

FLOWERS—White, small, in clusters on stalk 6–12 inches high.

LEAVES— Lobed and toothed. Hairy throughout, springing from the base.

2. MITREWORT; BISHOP'S CAP:
Mitélla diphýlla.

Saxifrage Family. Apr.—May.

Found in rich woods. Quebec to North Carolina, west to Minnesota and Missouri.

FLOWERS—White, tiny, on stalk 8–16 inches high.

LEAVES— Lower leaves coarsely toothed; 2 opposite leaves borne halfway up the flower stalk; hairy.

NAKED MITREWORT: *M. nùda.*

Resembles preceding. Flowers greenish and few. No leaves on flower stalk. Found in cool, mossy woods. Labrador to New York, Michigan, Minnesota and northward.

NOTES.

Digitized by Google



1. Foam-flower: False Mitrewort. **2. Mitrewort.**

Digitized by Google

1. EARLY SPRING, PLANTAIN-LEAVED, or MOUSE-EAR EVERLASTING; PUSSY'S TOES: White Plantain, and Ladies Tobacco. *Antennària plantaginifòlia.*

Composite Family. Apr.—June.

Found in dry fields and on hillsides. Common in U. S., Labrador to Gulf of Mexico, west to Nebraska.

FLOWER-HEADS—Whitish, in small, tight clusters terminating a white-woolly stem, 6–18 inches high.

LEAVES— Lower leaves broad near the tip, 3-ribbed, stalks nearly as long as the leaves; upper leaves narrow, downy.

2. EARLY SAXIFRAGE: *Saxífraga virginiénsis.*

Saxifrage Family. Apr.—June.

Found on dry, rocky hillsides. New Brunswick to Georgia, west to Minnesota and Tennessee.

FLOWERS—White, 5 divisions, small, in clusters terminating a downy stalk 4–10 inches high.

LEAVES— In a rosette at the base, toothed.

NOTES.

Digitized by Google

1. Early Everlasting. **2. Early Saxifrage.**

Digitized by Google

RED BANEBERRY; Cohosh: Herb-Christopher. *Actaèa rùbra.*

Crowfoot Family. Apr.—June.

Found in rich woods, an erect, bushy plant 1–2 feet high. Nova Scotia to Georgia and far west. Common northward.

FLOWERS—White, small, in short, thick, terminal cluster. Berries red, poisonous.

LEAVES— Compound, irregularly toothed and lobed.

WHITE BANEBERRY: *A. álba.*

Closely resembles the preceding except that the poisonous berries are white with a black eye. Common. Nova Scotia to Georgia and far west.

NOTES.

Digitized by Google



Red Baneberry.

Digitized by Google

SOLOMON'S SEAL: Hairy, or True, or Twin Flowered Solomon's Seal.

Polygónatum biflòrum.

Lily Family. Apr.—June.

Found in rich woods. A plant 1–3 feet high. New England to Florida, west to Michigan, Minnesota, Kansas and Texas.

FLOWERS—Greenish white, cylindrical, usually borne in pairs from the leaf angles. Berries blue-black.

LEAVES— Finely hairy beneath, set alternately on stem.

Rootstalk jointed, marked with scars of previous years' growths.

GREAT SOLOMON'S SEAL: *P. commutàtum.*

Found in meadows and along river-banks. A plant 2–7 feet high, resembles preceding, but leaves smooth throughout, partly clasping and flowers are in clusters of 2–8.

NOTES.

Digitized by Google

Solomon's Seal.

Digitized by Google

1. RUE ANEMONE: *Anemonélla thalictroìdes.*
Crowfoot Family. Apr.—June.
Found in thin woods, a small plant 5–9 inches high. New England to Minnesota, south to Tennessee.
FLOWERS—White, or pink tinged, 5–10 divisions, in clusters of 2–3.
LEAVES— In groups of 3, 3-lobed.
Roots tuberous.

2. WOOD ANEMONE: Wind-Flower:
Anemòne quinquefòlia.
Crowfoot Family. Apr.—June.
A plant 4–8 inches high. Found in open woods. Canada, south to Georgia, west to Rocky Mountains.
FLOWERS—White, or magenta tinged, solitary, 4–9 divisions.
LEAVES— In a whorl of 3–5, each leaf variously divided and toothed.

NOTES.

Digitized by Google

1. Rue Anemone. **2. Wood Anemone.**

Digitized by Google

1. WILD or FALSE SARSAPARILLA:
Aràlia nudicaúlis.

Ginseng Family. May—June.

A plant 8–12 inches high. Found in woods. New England to Dakota, south to mountains of Carolina.

FLOWERS—Greenish white, tiny, 3–7 clusters. Berries purple-black in clusters.

LEAVES— In 3 divisions, each consisting usually of 5 finely toothed leaflets.

Roots long, slender, aromatic.

2. DWARF GINSENG: Ground Nut.
Pànax trifòlium.

Ginseng Family. May—June.

A plant 4–8 inches tall. Found in woods. Canada to the mountains of Georgia.

FLOWERS—White, tiny, in small clusters. Berries yellow.

LEAVES— In 3 divisions, each consisting of 3–5 toothed leaflets.

Root spherical, aromatic.

NOTES.

Digitized by Google

1. Wild Sarsaparilla. **2. Dwarf Ginseng.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

ARROW ARUM: Green Arrow Arum.

Peltándra virgínica.

Arum Family. May—June.

A plant 1–3 feet high. Found growing in clumps in shallow water. Maine to Florida, west to Michigan and Missouri.

FLOWERS—Small, borne on club within a fleshy sheath; outer surface of sheath green. Berries green in close cluster.

LEAVES— Rich green, large, shaped like an arrow-head.

NOTES.

Digitized by Google



Arrow Arum.

Digitized by Google

1. WATER ARUM: Marsh Calla.

Cálla palústris.

Arum Family. May—June.

A plant 5–10 inches high. Found in cool bogs and swamps. Nova Scotia and southward to Virginia, westward to Minnesota, Iowa and Michigan.

FLOWERS—Tiny on short, thick club; outer envelope white above, greenish beneath, opening nearly flat. Berries red, in clusters.

LEAVES— Rich green, somewhat heart-shaped, from creeping rootstock.

2. MAIANTHEMUM: Canada Mayflower: Two-leaved Solomon's Seal: False Lily-of-the-Valley.

Maiánthemum canadénse.

Found in rich woods. Labrador to North Carolina, west to Dakota and Manitoba.

FLOWERS—White, tiny, in small clusters terminating a stem, 3–6 inches high. Berries brown-spotted, turning red in autumn.

LEAVES— 2–3 seated on stem.

NOTES.

Digitized by Google

1. Water Arum. 2. Maianthemum.

Digitized by Google

PAINTED TRILLIUM. *Trillium undulàtum.*
Lily Family. May—June.
Found in rich woods. Quebec and Ontario to Georgia, west to Wisconsin and Missouri.
FLOWERS—Solitary, 3 wavy-edged, white divisions marked with a crimson V at base.
LEAVES— Broad, in a whorl of 3 at the summit of stem, 8–15 inches high.

NODDING TRILLIUM: Nodding Wake-Robin. *T. cérnuum.*
FLOWERS—White, flower stem recurved so that the blossom hangs beneath the leaves.

LARGE-FLOWERED TRILLIUM: Large-flowered Wake-Robin or White Wood-Lily. *T. grandiflòrum.*
Found in same habitat, Western Vermont and Quebec to North Carolina, west to Minnesota and Missouri.
FLOWERS—White, turning pink with age, divisions often 2 inches long.

NOTES.

Digitized by Google



Painted Trillium.

Digitized by Google

1. STAR FLOWER: Star Anemone: Chickweed Wintergreen. *Trientàlis americàna.*
Primrose Family. May—June.
A plant 3–7 inches high. Found in moist woods. From mountains of Virginia and Illinois north to Labrador and Manitoba.
FLOWERS—White, on slender stalks, 6–7 pointed divisions.
LEAVES— 5–10 in a circle at summit of stem.

2. GOLDTHREAD: Canker-root. *Cóptis trifòlia.*
Crowfoot Family. May—July.
Found both in dry and wet woods. Labrador to Alaska, south to Michigan and mountains of North Carolina and Tennessee.
FLOWERS—White, 5–9 divisions, solitary, on stalk 3–6 inches high.
LEAVES— From the base, dark green, evergreen, finely toothed, divided into 3 leaflets. Rootstock wiry, yellow, bitter.

NOTES.

Digitized by Google



1. Star Flower. **2. Goldthread.**

Digitized by Google

FALSE SOLOMON'S SEAL: False Spikenard: Solomon's Zig-zag. *Smilacìna racemòsa.*

Lily Family. May—June.

A plant 1–3 feet high. Found in moist woods and borders of woods. Nova Scotia to Georgia, westward Minnesota, Arizona and British Columbia.

FLOWERS—White, tiny, in clusters terminating plant stem. Berries first brown-spotted, finally red.

LEAVES— Alternate, with very short stems if any.

STAR-FLOWERED SOLOMON'S SEAL *S. stellàta.*

Is a much smaller species.

FLOWERS—White, larger and fewer than those of the preceding. Berries black, or green, with black stripes.

LEAVES— Partly clasping.

NOTES.

Digitized by Google

False Solomon's Seal.

Digitized by Google

1. POISON SUMACH: *Rhús Vérnix.*
Cashew Family. June.

Shrub 6–15 feet high. Found on wet, low ground. Very poisonous. Western Maine to Ontario and southward.

FLOWERS—Small, whitish green in loose clusters from the angles of the leaves. Fruit small, dry, grayish white berries.

LEAVES— 7–13 leaflets without teeth. Leaf-stalks red.

2. POISON IVY: Three-Leaved Ivy: Mercury. *Rhús Toxicodéndron.*
Cashew Family. June—July.

A plant sometimes vine-like in habit, sometimes bushy. Found along roadsides, on fences and stone walls, tree-trunks, and running over the ground. Poisonous, and very common.

FLOWERS—Tiny, greenish white, in clusters. Fruit dry, grayish white berries.

LEAVES— In groups of 3 leaflets (Common Ivy, Virginia Creeper or Woodbine has 5), toothless or irregularly notched.

NOTES.

Digitized by Google

1. Poison Sumach. **2. Poison Ivy.**

Digitized by Google

BRISTLY SARSAPARILLA: *Aràlia hìspida.*
Ginseng Family. June—July.

Found in rocky woods. A hairy plant 2–3 feet high. Newfoundland to North Carolina, west to Indiana and Minnesota.

FLOWERS—Tiny, dull white, in clusters. Berries in clusters, bluish in effect, inedible.

LEAVES— Compound, toothed.

NOTES.

Digitized by Google

Bristly Sarsaparilla.

Digitized by Google

SHIN LEAF: *Pýrola ellíptica.*
Heath Family. June—July.
Found in woods. Eastern United States and Quebec to Maryland, west to Rocky mountains.
FLOWERS—Greenish white, 5 divisions, fragrant, on stalk 5–10 inches high.
LEAVES— From the base, thin, exceeding their stalks in length.

ROUND-LEAVED PYROLA: *P. americàna.*
FLOWERS—Similar to the preceding.
LEAVES— Nearly round, thick, shining green.

ONE-FLOWERED PYROLA: *Monèses uniflòra.*
Bears a single white or pink flower.

NOTES.

Digitized by Google



Shin Leaf.

Digitized by Google

1. RED-BERRIED ELDER:
Sambùcus racemòsa.
Honeysuckle Family. June—July.
Found on borders of woods. A shrub 2–12 feet high. New Foundland to British Columbia, south to Georgia, Iowa and Colorado.
FLOWERS—Dull white, in large cluster, followed by red berries.
LEAVES— 5–7 finely-toothed leaflets.

COMMON ELDER: Elderberry: American, Sweet, or Black-berried Elder.
S. canadensis.
FLOWERS—White, in flat clusters; fruit purple-black.

2. SWEET PEPPERBUSH; White Alder: Alder-leaved Clethra. *Clèthra alnifòlia.*
Heath Family. July—Aug.
Shrub 3–10 feet high. Found in low woodlands, thickets, chiefly near the Atlantic coast.
FLOWERS—White, in narrow clusters, fragrant.
LEAVES— Alternate, finely toothed, especially toward the tips.

NOTES.

Digitized by Google

1. Red-berried Elder. **2. Sweet Pepperbush.**

Digitized by Google

1. REIN ORCHIS: *Habenària bracteàta.*
Orchis Family. June—Aug.
Found in damp woods. Nova Scotia to Alaska, south to Washington, Minnesota and Pennsylvania.
FLOWERS—Greenish, small, in spike terminating stem 6–20 inches high.
LEAVES— Long and broad. Long, narrow bracts or leaflets spring from bases of flowers.

2. REIN ORCHIS: *Habenària hyperbòrea.*
Orchis Family. June—Aug.
Found in wet woods. Newfoundland and Alaska, south to Pennsylvania and Nebraska.
FLOWERS—Greenish, in dense, narrow spike terminating stem 8–30 inches high.
LEAVES— Oblong to lance-shaped.

NOTES.

Digitized by Google

1. Rein Orchis. 2. Rein Orchis.

Digitized by Google

1. THIMBLEWEED: TALL or SUMMER ANEMONE. *Anemòne virginiàna.*
Crowfoot Family. June—Aug.
Found along roadsides and borders of woodlands. Maine to Minnesota and southward.
FLOWERS—White or greenish, 5 divisions, terminating stalks 2–3 feet high. Seeds form thimble-shaped burrs.
LEAVES— 3 in a whorl, 3-parted, divisions variously lobed and toothed. Later flowers bear 2 divided leaves midway of flower stalks.

2. FIELD CHICKWEED: *Cerástium arvénse.*
Pink Family. May—July.
Found in dry, rocky land. Common.
FLOWERS—White, 5 divisions, divisions 2-lobed, terminating stems 4–10 inches high.
LEAVES— Small, narrow.

NOTES.

Digitized by Google

1. Thimbleweed. **2. Field Chickweed.**

Digitized by Google

1. INDIAN PIPE; Corpse Plant: Ice-plant: Ghost-flower. *Monótropa uniflòra.*
Heath Family. June—Aug.
Found in moist, heavily-shaded woods. Throughout North America, except far south.
FLOWERS—Waxy white, solitary, nodding, terminating stem 4–10 inches high.
LEAVES— None. Plant stem colorless, turning black with age, bearing numerous scales, one or more stems arising from a mass of brittle, fibrous roots.

2. WILD LEEK: *Allium tricóccum.*
Lily Family. June—July.
Found in rich woods. New Brunswick to Minnesota, south to North Carolina.
FLOWERS—Greenish white, clustered at the end of stalk 4–15 inches high.
LEAVES— Two or three leaves 8–10 inches long, which wither before the flowers bloom, arise from a pointed bulb; strongly onion scented.

Digitized by Google

1. Indian Pipe. **2. Wild Leek.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

1. BLADDER CAMPION: *Silène latifòlia.*
Pink Family. June—Aug.
Found in fields, roadsides, waste places. Canada south to New Jersey, Illinois and Iowa. Naturalized from Europe.
FLOWERS—White, 5 divisions, divisions deeply 2-lobed, flower cup much inflated; terminating branches 8–18 inches high.
LEAVES— Opposite, smooth.

2. WHITE CAMPION: Evening Lychnis.
Lýchnis álba.
Pink Family. July—Sept.
Plant 1–2 feet high. Found in waste places. Adventive from Old World.
FLOWERS—White or pink, fragrant, opening in the evening; divisions 5, deeply cleft, sticky and hairy beneath the flower. Small clusters terminate branches.
LEAVES— Opposite, leaves and stems hairy.

Digitized by Google

1. Bladder Campion. **2. White Campion.**

Digitized by Google

1. MUSK MALLOW: *Málva moschàta.*
Mallow Family. June—Sept.
Found by roadsides and in waste places. Canada, Eastern, Middle and Southern States. Naturalized from Europe. Escaped from cultivation.
FLOWERS—White or pink, 5 divisions, in small cluster terminating stem, 1–2 feet high.
LEAVES— Deeply cut and lobed, faintly musk-scented.

2. COMMON MALLOW; Cheeses.
Málva rotundifòlia.
Mallow Family. June—Oct.
Found in waste places. Throughout United States. Naturalized from Europe.
FLOWERS—White or pinkish, small, on long stalks from leaf angles.
LEAVES— Slightly lobed, long stemmed. Plant stem creeping, 4–10 inches long.

Digitized by Google



1. Musk Mallow. **2. Common Mallow.**

Digitized by Google

1. CHAMOMILE: Mayweed: Dog-fennel: Dill-weed: Pig-sty Daisy: Dog's or Fetid Chamomile. *Anthemis Cótula.*

Composite Family. June—Oct.

A plant 8–20 inches high. Found by roadsides and in dry waste land. Widely distributed throughout North America.

FLOWER-HEADS—Resemble small daisies, the white rays 3-toothed on the tip.

LEAVES— Finely dissected, ill-scented.

2. BUNCHBERRY: Low or Dwarf Cornel. *Córnus canadénsis.*

Dogwood Family. June—July.

Found in woods. Plant 4–8 inches high. Labrador to Alaska, south to West Virginia and Minnesota.

FLOWERS—Dull white, tiny, in clusters surrounded by 4 white leaflets simulating petals. Berries red in close cluster, edible.

LEAVES— Smooth, oval, pointed, 4–6 in a whorl.

Digitized by Google

Chamomile. **2. Bunchberry.**

Digitized by Google

DAISY FLEABANE: *Erígeron ramòsus.*
Composite Family. June—Oct.

A plant 1–2 feet high. Found in fields, waste lands, roadsides. Nova Scotia to Virginia, and west to Missouri.

FLOWER-HEADS—About ½ inch across, with large, green-yellow disk, rays white, narrow.

LEAVES— Narrow, nearly or quite toothless, lower leaves broader at the tip.

DAISY FLEABANE; SWEET SCABIOUS: *E. ánnuus.*

Similar to the preceding, with broader, coarsely-toothed leaves. Rays white or pale lilac.

NOTES.

Digitized by Google



Daisy Fleabane.

Digitized by Google

YARROW; MILFOIL: Old Man's Pepper; Nose-bleed. *Achillèa Millefòlium.*

Composite Family. June—Oct.

Found on roadsides and in waste places. Common throughout North America.

FLOWER-HEADS—Grayish white, sometimes crimson, small, in flat-topped cluster terminating stem, 1–2 feet high.

LEAVES— Finely dissected, aromatic.

NOTES.

Digitized by Google

Yarrow.

Digitized by Google

VIRGIN'S BOWER: Traveller's Joy: Old Man's Beard: Devil's Yarn.

Clématis virginiàna.

Crowfoot Family. July—Aug.

A vine 6–12 feet long. Found in rich, damp soil, climbing over fences and shrubs. From Canada south to Georgia and Kansas.

FLOWERS—White or greenish, 4 divisions, in loose clusters from the leaf angles. Seeds in plumy masses.

LEAVES— Opposite, consisting each of 3 coarsely-toothed leaflets.

NOTES.

Digitized by Google

Virgin's Bower.

Digitized by Google

1. RAGGED FRINGED ORCHIS: Fringed Green Orchis. *Habenària lácera.*
Orchis Family. July—Aug.
Found in swamps and wet woods from Newfoundland to Minnesota, south to Missouri and Alabama.
FLOWERS—Greenish or dingy white, with lower lip cut to thread-like fineness; borne in spike terminating stem, 10–20 inches high.
LEAVES— Long and narrow, or oblong.

2. RATTLESNAKE PLANTAIN: Downy Rattlesnake Plantain. *Epipáctis pubéscens.*
Orchis Family. July—Aug.
Found in dry woods. New England to Florida, west to the Mississippi.
FLOWERS—Small, white, numerous on downy stalk 6–18 inches high.
LEAVES— Mottled with light and dark green.

Digitized by Google

1. Ragged Fringed Orchis. **2. Rattlesnake Plantain.**

Digitized by Google

GREATER GREEN ORCHIS: Large Round-leaved Orchis. *Habenària orbiculàta.*
Orchis Family. July—Aug.
Found in rich woods. Labrador to Alaska, south to South Carolina, Minnesota, and Washington.
FLOWERS—Greenish white, on stalk 1–2 feet high.
LEAVES— 2 large, broad leaves, shining green above, silvery beneath, lying flat on the ground. One or more tiny leaflets borne on the flower stalk.

HOOKER'S ORCHIS: *H. Hookèri.*
Similar to the preceding. Flowers yellowish green. No leaflets on flower stalk.

NOTES.

Digitized by Google

Greater Green Orchis.

Digitized by Google

1. ROUND-LEAVED SUNDEW: Dew-plant.
Drósera rotundifòlia.
Sundew Family. July—Aug.
Found in bogs and sandy marshes. Labrador to the Gulf of Mexico and westward. From Alaska to California.
FLOWERS—Small, white, opening one at a time, on stalk 4–10 inches high.
LEAVES— Grow in a rosette, round, borne on long stems, and covered with red hairs.

LONG-LEAVED SUNDEW: *D. longifòlia.*
Leaves widest at the tip, tapering toward the stems. Leaf stems without hairs.

2. LADIES' TRESSES:
Spiránthes cérnua.
Orchis Family. Aug.—Oct.
Found in low meadows and swamps. Newfoundland to Georgia, Minnesota and Nebraska.
FLOWERS—Small, white or yellowish, arranged spirally on stem 6–12 inches high.
LEAVES— Long and narrow, grass-like.

SLENDER LADIES' TRESSES:
S. grácilis.
Found in pastures and on dry hillsides. Bears flowers on one side of the stem, or spirally. Basal leaves wither before flowers bloom. Roots tuberous.

Digitized by Google



1. Round-leaved Sundew. 2. Ladies' Tresses.

Digitized by Google

THOROUGHWORT; BONESET: Ague-weed: Indian Sage. *Eupatòrium perfoliàtum.*
Composite Family. July—Sept.
Found in low, wet ground. From Nebraska, Manitoba, New Brunswick, south to the Gulf of Mexico.
FLOWER-HEADS—Small, dull white, in large, flat-topped cluster terminating stem 2–5 feet high.
LEAVES— Opposite, often united at the base around the stem, wrinkled, toothed. Plant stem and under surface of leaves downy.

WHITE SNAKEROOT: White or Indian Sanicle: Deerwort Boneset. *E. urticaefòlium.*
Found in rich moist woods. A plant 1–4 feet high.
FLOWER-HEADS—White, resembling the garden Ageratum, in spreading cluster.
LEAVES— Nearly heart-shaped, coarsely toothed, long-stemmed.

NOTES.

Digitized by Google



Thoroughwort.

Digitized by Google

TURTLEHEAD; BALMONY: Snake-head: Shell-flower: Cod-head. *Chelòne glàbra.*
Figwort Family. July—Sept.
Found in wet ground. Newfoundland to Florida and Texas, west to Mississippi and Minnesota.
FLOWERS—White or tinged with pink, terminating stem 1–3 feet high.
LEAVES— Opposite, toothed.

NOTES.

Digitized by Google



Turtlehead.

Digitized by Google

1. ARROW-HEAD: *Sagittària latifòlia.*
Water Plantain Family. July—Sept.

Found in shallow water. From Mexico northward throughout North America.

FLOWERS—White, divisions 3, in whorls of 3, on stalk 6 inches to 4 feet high.

LEAVES— Arrow-head-shaped, exceedingly variable, from short and broad to long and narrow.

2. ROUGH BEDSTRAW: *Gàlium asprèllum.*
Madder Family. June—Aug.

A plant 2–4 feet tall. Found in rich, moist ground. Eastern half of United States and Canada.

FLOWERS—White, tiny, in airy clusters. Flower stems 2–3 times forked.

LEAVES— In whorls of 6 on main stems, prickly-rough on edge and rib. Plant stems rough backwards with hooked prickles.

Digitized by Google



1. Arrow-head. **2. Rough Bedstraw.**

Digitized by Google

NOTES.

Digitized by Google

Digitized by Google

WILD BALSAM-APPLE: *Echinocýstis lobàta.*
Gourd Family. July—Sept.

A rapid climbing vine 10–20 feet long. Found in rich, moist soil, beside streams and in waste places. New Brunswick to Kentucky, west to Manitoba and Texas.

FLOWERS—Greenish white, small, in loose clusters from the leaf angles.

LEAVES— Deeply and sharply 5-lobed.

NOTES.

Digitized by Google

Wild Balsam-apple.

Digitized by Google

TALL MEADOW RUE:

Thalictrum polygamum

Crowfoot Family. July—Sept.

Found in swamps and low meadows. Newfoundland to Ohio and southward.

FLOWERS—White or greenish, in plume-like cluster terminating stem 3–10 feet high.

LEAVES— Compound, the leaflets small, lustreless, lobed.

NOTES.

Digitized by Google

Tall Meadow Rue.

Digitized by Google

GRASS OF PARNASSUS: Carolina Grass of Parnassus. *Parnássia caroliniàna.*

Saxifrage Family. July—Sept.

Found in swamps and wet meadows. New Brunswick to Florida, west to Iowa.

FLOWERS—Creamy white with greenish veins, divisions 5, on slender stalks 8–20 inches high.

LEAVES— From the base, somewhat heart-shaped, a single leaf clasping the flower stalks.

NOTES.

Digitized by Google

Grass of Parnassus.

Digitized by Google

WATER PLANTAIN:

Alisma Plantàgo-aquática.

Water Plantain Family. July—Sept.

Found in shallow water. North America, Europe, Asia.

FLOWERS—Small, white, in loose, symmetrical cluster, on stalk 1–3 feet high.

LEAVES— From the base, variable in shape, broad or narrow.

NOTES.

Digitized by Google



Water Plantain.

Digitized by Google

1. COMMON DODDER: Strangle-weed: Love-vine: Angel's Hair. *Cúscuta Gronòvii.*
Convolvulus Family. July—Sept.
Found in moist soil. Nova Scotia and Minnesota south to Gulf of Mexico.
FLOWERS—Tiny, dull white, 5-lobed, in clusters.
LEAVES— None. Stems long, wiry, yellow or orange, twining over vegetation.

2. WOOD SORREL: White or True Wood-sorrel: Alleluia: *Oxalis Acetosélla.*
Wood Sorrel Family. May—July.
A plant 3–4 inches tall. Found in cold, damp woods. Nova Scotia and Manitoba, south to North Carolina.
FLOWERS—White or pinkish, veined with deep pink lines, 5 divisions.
LEAVES— From the base, composed of 3 heart-shaped leaflets.

NOTES.

Digitized by Google

1. Dodder. **2. Wood Sorrel.**

Digitized by Google

1. PEARLY, or LARGE-FLOWERED, EVERLASTING: Immortelle: Silver-leaf: Cotton-weed: Moonshine: None-so-pretty.

Anáphalis margaritàcea.

Composite Family. July—Sept.

Found in dry fields, hillsides. North Carolina to Kansas and California, also far north.

FLOWER-HEADS—Center yellow surrounded by white scales, the cluster terminating a white-woolly stem 1–3 feet high.

LEAVES— Long, narrow, white-woolly.

2. WHITE MELILOT; SWEET CLOVER: Bokhara or Tree Clover: Honey Lotus.

Melilòtus álba.

Pulse Family. June—Aug.

Found by roadsides and in waste places. Naturalized from Europe: widespread in United States.

FLOWERS—White, small, fragrant, in slender clusters terminating branches 3–10 feet high.

LEAVES— Compound, consisting of 3 finely-toothed leaflets.

Digitized by Google

1. Pearly Everlasting. **2. White Melilot.**

Digitized by Google

WILD CARROT; QUEEN ANNE'S LACE:
Bird's Nest. *Daúcus Caròta.*
Parsley Family. July—Sept.

A plant 2–3 feet high. Found in fields, roadsides, waste places. Naturalized from Europe. Eastern half of United States and Canada.

FLOWERS—White, tiny, in broad, rather flat cluster, curling up later, resembling a bird's nest.

LEAVES— Finely dissected.

NOTES.

Digitized by Google

Wild Carrot.

Digitized by Google

BUTTONBUSH: Globe-flower: Honey-balls: Button-ball Shrub: River-bush.

Cephalánthus occidentàlis.

Madder Family. **July—Sept.**

A shrub 3–12 feet high. Found in wet ground and beside ponds and streams. New Brunswick to Florida and Cuba, westward to Arizona and California.

Flowers—White, small, clustered in round heads.

Leaves— Opposite, or in whorls of 3.

NOTES.

Digitized by Google



Buttonbush.

Digitized by Google

POKEWEED; PIGEON BERRY: Scoke: Inkberry: Garget. *Phytolácca decándra.*

Pokeweed Family. July—Sept.

A plant 4–10 feet tall. Found by roadsides, in clearings, waste places. Maine and Ontario to Florida and Texas.

FLOWERS—Small, with 5 white divisions, pink tinted outside, and a conspicuous green center; borne in loose spikes. Berries dark purple in long drooping clusters. Ripe August—October.

LEAVES— Oblong to lance-shaped, tapering at both ends.

Plant stems branching, reddening toward fall. Root large, poisonous.

NOTES.

Digitized by Google

Pokeweed.

Digitized by Google

FLOWERING MOSS; PYXIE: Pine-barren Beauty. *Pyxidanthèra barbulàta.*

Diapensia Family. Mch.—May.

A low plant, growing in mat-like patches, creeping prostrate branches 6–10 inches long. Found in dry, sandy soil, pine barrens. New Jersey to North Carolina.

FLOWERS—White, often pink, about 1/4 inch across, 5-lobed, numerous.

LEAVES— Very small and narrow, seated on stem, crowded toward the ends of the branches, moss-like in appearance.

MAYFLOWER; TRAILING ARBUTUS: Ground Laurel. *Epigaèa rèpens.*

Heath Family. Mch.—May.

Found in woodlands or mossy, rocky places. Newfoundland to Florida, west to Kentucky, Minnesota and the Northwest.

FLOWERS—White or pink tinted, 5-lobed, fragrant, in clusters at the ends of the woody, creeping branches 6–10 inches long.

LEAVES— Oval, rounded at the base, old leaves more or less rusty spotted.

Digitized by Google

Digitized by Google

WHITE ADDER'S TONGUE:
Erythrònium álbidum.
Lily Family. Mch.—May.
Found in rich soil. Common west as far as Texas, Arkansas and Minnesota. Somewhat rare in Eastern States, New York to New Jersey.
FLOWERS—Pinkish-white, yellow at the center, purplish-tinged outside, 6 strongly recurved divisions, terminating stalks 5–8 inches high.
LEAVES— 2, from the base, smooth, more or less mottled, from solid bulb.

TWINLEAF: Rheumatism Root.
Jeffersònia diphýlla.
Barberry Family. Apr.—May.
Found in rich, shady woods. New York to Wisconsin, south to Tennessee.
FLOWERS—White, 8 divisions, 1 inch broad, solitary, on stalk about 7 inches high, twice as tall in seed time.
LEAVES— From the root, long-stemmed, parted into 2 divisions.

Digitized by Google

MAY APPLE; MANDRAKE: Hog Apple: Wild Lemon. *Podophýllum peltàtum.*
Barberry Family. Apr.—May.

A plant 12–18 inches high. Found in rich, moist woods. Western Quebec and western New England to Minnesota and southward.

FLOWERS—White, 6–9 divisions, solitary, unpleasantly scented. Fruit, large, edible berry.

LEAVES— Those of the flowerless stems terminated by a large, rounded, 7–9-lobed leaf supported by a stem in the middle; those of the flowering stem 2, much lobed, with the nodding flower in the fork.

SWEET WHITE VIOLET: *Viola pállens.*
Violet Family. Apr.–May.

Plant 3–5 inches tall. Found in moist, wooded places. Common.

FLOWERS—White with purple veins, side divisions usually bearing a small tuft of hairs, slightly fragrant.

LEAVES— Round heart-shaped, slightly scalloped.

Digitized by Google

LANCE-LEAVED VIOLET: *Viola lanceolàta.*

Violet Family. Apr.—June.

Found in moist ground. Nova Scotia to Minnesota and southward.

FLOWERS—White, purple veined, side divisions beardless.

LEAVES— Long, narrow, indistinctly scalloped, tapering into a long stem.

CANADA VIOLET: *Viola canadénsis.*

Violet Family. May—July.

Found in woodlands, very widely-distributed, mountainous and hilly country chiefly. Easily distinguished from other white violets by its tall leafy stem 6–24 inches high.

FLOWERS—White or pale lavender, violet tinged on the outside, side divisions bearded, lower one striped with fine dark lines.

LEAVES— Heart-shaped, pointed, toothed.

NOTES.

Digitized by Google

WATER CRESS:

Radícula Nastúrtium-aquáticum.

Mustard Family. Apr.—Aug.

Found in brooks, ditches. Naturalized from Europe. Common.

FLOWERS—White, 4 divisions, small, in terminal clusters.

LEAVES— Compound, with 3–11 rounded leaflets. Stems spreading and rooting, 4–10 inches high.

CREEPING SNOWBERRY:

Chiógenes hispídula.

Heath Family. May—June.

A low, creeping plant. Found in mossy bogs and woods. Canada, south to Minnesota, Michigan and North Carolina.

FLOWERS—White, very small, bell-shaped, 4-lobed, solitary, from the leaf angles. Berries white.

LEAVES— Evergreen, small, with an abrupt point. Leaves and berries aromatic.

NOTES.

Digitized by Google

NEW JERSEY TEA; RED-ROOT:
Ceanòthus americànus.
Buckthorn Family. May—July.
A shrub 1–4 feet high. Found in dry, open woods. Gulf of Mexico to Ontario.
FLOWERS—White, small, in dense, oblong, terminal clusters.
LEAVES— Long egg-shaped, finely toothed, conspicuously 3-ribbed. Root reddish.

BLACK SNAKEROOT; BLACK COHOSH:
Tall Bugbane. *Cimicífuga racemòsa.*
Crowfoot Family. June—July.
Found in rich woods, hillsides. Southern New England and Wisconsin south to Georgia.
FLOWERS—White, ill-scented, fuzzy, borne on a long, slender spike terminating stem 3–8 feet high.
LEAVES— Compound, sharply toothed.

CLAMMY AZALEA; WHITE SWAMP HONEYSUCKLE: *Rhododéndron viscòsum.*
Heath Family. June—July.
A shrub 3–7 feet high, resembling the earlier-blooming Pink Azalea. Found in swamps, generally near the coast. Maine, south and west to Ohio and Arkansas.
FLOWERS—White or pink-tinged, 5-lobed; tube longer than the lobes, sticky-hairy outside.
LEAVES— Blunt lance-shaped, margins and midrib bristly.

Digitized by Google

SHOWY LADY'S SLIPPER: *Cypripèdium hirsùtum.*

Orchis Family. June—July.

Found in wet, mossy woods, swamps. Newfoundland to Georgia, west to Mississippi.

FLOWERS—Pouch white, more or less stained with crimson-magenta, similar in shape to the Stemless Lady's Slipper, but differs in the pouch being open but not divided; other divisions of the flower white, wide, not longer than the pouch.

LEAVES— Broad, pointed, many ribbed. Plant stem leafy 1–2 feet high.

LIZARD'S TAIL: *Saurùrus cérnuus.*

Pepper Family. June—Aug.

A plant 2–5 feet high. Found in swamps, shallow water. Rhode Island to Gulf of Mexico, west to Minnesota and Texas.

FLOWERS—White, fragrant, very small, crowded on slender spikes, nodding at the tip.

LEAVES— Heart-shaped, converging ribs, dark green, on stout stems.

Plant-stem jointed, sparingly branched, leafy.

NOTES.

Digitized by Google

POKE or TALL MILKWEED:
Asclèpias phytolaccoìdes.
Milkweed Family. June—Aug.
A plant 3–6 feet high. Found in woodland borders, thickets from Maine to Georgia and far west.
FLOWERS—Similar in structure to the Common Milkweed, white and greenish, or magenta tinged, drooping, in a loose cluster.
LEAVES— Rather broad, pointed at each end.

WATER HEMLOCK; SPOTTED COW-BANE: Musquash Root: Beaver Poison.
Cicùta maculàta.
Parsley Family. June—Aug.
Found in wet meadows, swampy places. A deadly poison contained in its fleshy roots. New Brunswick to Virginia and westward.
FLOWERS—White, tiny, in thin, flat clusters terminating branches, 3–6 feet high.
LEAVES— Compound, leaflets toothed. Plant stem marked with purplish lines.

WATER PARSNIP: Hemlock Water Parsnip.
Sìum cicutaefòlium.
Is a similar species growing in wet places and also poisonous. It may be fóund from the Atlantic to the Pacific.
FLOWERS—White, in broad clusters.
LEAVES— Compound 7–15 sharply toothed leaflets, but finely dissected when growing under water.

Digitized by Google

DALIBARDA: Creeping Dalibarda.

Dalibárda rèpens.

Rose Family. June—Sept.

Found in woodlands. Nova Scotia to Pennsylvania, west to the Mississippi.

FLOWERS—White, solitary, 5 divisions, on stalks 2–5 inches high.

LEAVES— From the base, heart-shaped, scallop-toothed; leaves and stems finely hairy.

WATER LILY: Pond Lily: Water Nymph: Sweet-scented Water Lily.

Castàlia odoràta.

Water Lily Family. June—Sept.

Found in shallow water. Common. Nova Scotia to the Gulf of Mexico, west to Mississippi.

FLOWERS—Pure white, sometimes pink tinged, center yellow, divisions many. Flower 3–5 inches across, floating, very fragrant.

LEAVES— Floating, mostly oblong in shape, cleft at the base, on long stems.

NOTES.

Digitized by Google

WHITE FRINGED ORCHIS:
Habenària blephariglóttis.
Orchis Family. July—Aug.
Found in bogs, swamps. Eastern Canada to Michigan, south to Mississippi.
FLOWERS—Similar in habit to the Yellow Fringed Orchis, white, fragrant, lip fringed, in spike terminating stem 1–2 feet high.
LEAVES— Lance-shaped, parallel veined, clasping stem.

PRAIRIE WHITE FRINGED ORCHIS:
H. leucophaèa.
Is similar to the preceding, differing chiefly in having larger white or greenish flowers, lip divided into wedge-shaped segments, deeply fringed. It is more common west, being found in moist, open ground from Western New York to Minnesota and Arkansas.

WINTERGREEN; CHECKERBERRY: Partridge-berry: Deer, Box, or Spice-berry: Mountain or Ground Tea. *Gaulthèria procúmbens.*
Heath Family. July—Aug.
Found in woodlands. Newfoundland to Northern Georgia, west to Minnesota, Michgan and Manitoba and far north.
FLOWERS—White, small, somewhat bell-shaped, nodding, from the leaf angles. Berries red, edible.
LEAVES— Dark green, evergreen, slightly toothed, aromatic, clustered at the top of stem, 2–5 inches high.

Digitized by Google

CULVER'S ROOT: Culver's Physic:
Verónica virgínica.
Figwort Family. July—Aug.
An erect plant with straight, usually unbranched stem 2–7 feet high. Found in moist woods, meadows. Western Massachusetts and Connecticut to Manitoba and southward.
FLOWERS—White or pale lavender, in dense, terminal spikes 3–6 inches long.
LEAVES— 4–7, in whorls, lance-shaped, pointed, finely toothed.

JAMESTOWN or JIMSON WEED: Thorn-apple: Stramonium: Devil's Trumpet.
Datùra Stramònium.
Nightshade Family. July—Sept.
A plant 1–5 feet tall. Found in fields, waste land. Nova Scotia to Gulf, west beyond the Mississippi Naturalized from Asia.
FLOWERS—White, about 4 inches long, trumpet-shaped, with 5 sharply-pointed lobes. Fruit large, egg-shaped, covered with spines.
LEAVES— Coarsely toothed and angled, rank-smelling.

ASTERS—See page 304.

NOTES.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

II

YELLOW AND ORANGE

Digitized by Google

YELLOW AND ORANGE

YELLOW ADDER'S-TONGUE; DOG'S-TOOTH VIOLET: Trout Lily.

Erythrònium americànum.

Lily Family. Apr.—May.

Found in moist woods, brooksides. New Brunswick to Florida, west to Mississippi, Minnesota and Arkansas.

FLOWERS—Lily-like, 6 divisions, yellow, more or less tinged with brown-purple on the outside, borne on stalk 6–12 inches high.

LEAVES— 2 from the base, green, or mottled with brown-purple.

NOTES.

Digitized by Google

Yellow Adder's Tongue.

Digitized by Google

MARSH MARIGOLD: Meadow-gowan: American Cowslip. ***Caltha palustris.***

Crowfoot Family. **Apr.—May.**

A plant 8–24 inches high. Found in low, wet meadows. Newfoundland to South Carolina, Tennessee, and Nebraska.

Flowers—Yellow, 5–9 divisions.

Leaves— Mostly from the root, round, kidney or heart-shaped.

NOTES.

Digitized by Google

Marsh Marigold.

Digitized by Google

1. STAR GRASS: *Hypóxis hirsùta.*
Amaryllis Family. Apr.—July.
Found in dry, open woods, meadows. Maine to Gulf of Mexico and west to Minnesota, Texas, Kansas.
FLOWERS—Yellow inside, greenish and hairy outside, 6 divisions.
LEAVES— Grass-like, 2–6 inches high.

2. CINQUEFOIL: Common Cinquefoil: Five-Finger. *Potentílla canadénsis.*
Rose Family. Apr.—Aug.
Found in dry fields, roadsides, waste places. Quebec to Georgia, west to Minnesota, Iowa.
FLOWERS—Yellow, 5 divisions, growing singly on slender stalks from leaf angles.
LEAVES— Composed of 5 toothed leaflets. Plant stems wiry, prostrate, 6–20 inches long.

NOTES.

Digitized by Google

1. Star Grass. **2. Cinquefoil.**

Digitized by Google

1. GOLDEN ALEXANDERS: Early Meadow Parsnip. *Zizia aúrea.*

Parsley Family. May—June.

Found in meadows and river-banks. From New Brunswick and Dakota to the Gulf States.

FLOWERS—Yellow, tiny, in thin, flat clusters terminating branches, 8–30 inches high.

LEAVES— Compound, generally in 3 divisions, each composed of 3–7 toothed leaflets.

2. WILD PARSNIP; FIELD PARSNIP; Madnep: Tank. *Pastinàca sativa.*

Parsley Family. June—Sept.

A plant 2–5 feet tall, with noticeably grooved stem, found in roadsides, waste places. Common throughout United States and Canada. Naturalized from Europe.

FLOWERS—Dull yellow, tiny, forming flat clusters terminating branches, 2–5 feet high.

LEAVES— Compound, composed of many toothed leaflets.

NOTES.

Digitized by Google

1. Golden Alexanders. **2. Wild Parsnip.**

Digitized by Google

CLINTONIA; YELLOW CLINTONIA:

Clintònia boreàlis.

Lily Family. May—June.

Found in moist woods. Labrador to North Carolina, west to Manitoba and Minnesota.

FLOWERS—Greenish yellow, 6 divisions, lily-like, 3–6 terminating stalk 6–15 inches high. Berries pure blue.

LEAVES— Usually 3, from the base, oval or oblong.

NOTES.

Digitized by Google

Clintonia.

Digitized by Google

INDIAN CUCUMBER-ROOT:

Medèola virginiàna.

Lily Family. May—June.

A plant 1–3 feet high, found in rich woods. New Brunswick to Minnesota, south to Gulf States.

FLOWERS—Greenish yellow, few at summit of stem. Berries purple-black.

LEAVES— In 2 whorls; lower whorl of 5–9 leaves, upper whorl of 3–5 smaller leaves. Rootstock white, with slight cucumber flavor.

NOTES.

Digitized by Google



Indian Cucumber-root.

Digitized by Google

1. OAKESIA: *Oakèsia sessilifòlia.*
Lily Family. May—June.
A plant 6–12 inches tall, found in rich woods. New England to Florida, west to Minnesota, Nebraska and Arkansas.
FLOWERS—Pale yellow, 6 divisions, long and slender. Seed capsule 3-sided, resembling a beechnut.
LEAVES— Partly clasping, whitish beneath.

2. LARGE-FLOWERED BELLWORT: *Uvulària grandiflòra.*
Lily Family. Apr.—June.
Found in rich woods. Western New England to Minnesota and Kansas, south to Georgia.
FLOWERS—Larger and more irregular than those of the preceding, pale or greenish yellow, inner surface smooth.
LEAVES— Leaves appear as if perforated by the plant stem.

3. BELLWORT: *U. perfoliàta.*
Resembles the preceding but may be distinguished by the rough inner surface of the flower. Found from Massachusetts to Ontario, Dakota, and southwest.

NOTES.

Digitized by Google



1. Oakesia. 2. Large-flowered Bellwort. 3. Bellwort.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

YELLOW LADY'S SLIPPER: Large Yellow Lady's Slipper: Yellow Moccasin Flower: Whippoorwill's Shoe.

Cypripèdium parviflòrum Var. pubéscens.

Orchis Family. May—July.

Found in moist or boggy woods. Nova Scotia to Alabama, west to Minnesota, E. Kansas and Nebraska.

FLOWERS—Solitary, borne at the top of leafy stem 1–2 feet high; pouch golden yellow; other divisions of flower lined and marked with madder-purple.

LEAVES— Parallel veined, clasping stem.

SMALL YELLOW LADY'S SLIPPER: *C. parviflòrum.*

Similar but smaller than the preceding, with similar range. Intergrading forms frequent.

NOTES.

Digitized by Google

Yellow Lady's Slipper.

Digitized by Google

GOLDEN RAGWORT; Groundsel: Squaw-weed.
Senècio aúreus.

Composite Family. May—July.

Found in wet meadows. Newfoundland and Ontarió to Gulf of Mexico, west to Missouri.

FLOWER-HEADS—Golden yellow, in loose clusters terminating stem, 12–30 inches high.

LEAVES— Those from the root round or heart-shaped, long-stemmed, toothed; stem leaves variable, deeply lobed, uppermost clasping.

NOTES.

Digitized by Google



Golden Ragwort.

Digitized by Google

CARRION-FLOWER: *Smilax herbàcea.*
Lily Family. May—July.

A handsome vine growing 4–15 feet long, found in moist soil. Canada to Gulf States, west to Nebraska.

FLOWERS—Greenish yellow, in inconspicuous clusters, ill-scented. Berries at first green, finally blue-black.

LEAVES— Somewhat heart-shaped, parallel veined.

NOTES.

Digitized by Google

Carrion-flower.

Digitized by Google

YELLOW POND LILY; COW LILY: Large Yellow Pond or Water Lily: Spatter-dock.

Nymphæa ádvena.

Water Lily Family. May—Sept.

Found in shallow, stagnant water. Nova Scotia to Gulf States, west to Rocky Mountains.

FLOWERS—Yellow, more or less tinged with green and purple, 6 unequal divisions; flowers borne erect on thick stems.

LEAVES— Large, deeply cleft at the base, floating.

NOTES.

Digitized by Google



Yellow Pond Lily.

Digitized by Google

WILD YELLOW LILY: Meadow or Field Lily: Canada Lily. *Lílium canadénse.*
Lily Family. June—July.
Found in wet meadows and swamps. Nova Scotia to Georgia, west beyond Mississippi, Minnesota and Missouri.
FLOWERS—Yellow to orange, with purple-brown spots inside, terminating stems 2–5 feet high.
LEAVES— Usually in whorls of 4–10.

TURK'S-CAP LILY: Turban Lily. *L. supérbum.*
Flowers orange-yellow, half drooping, divisions more reflexed than those of the preceding, thickly spotted. Upper leaves arranged alternately, lower leaves in whorls.

NOTES.

Digitized by Google

Wild Yellow Lily.

Digitized by Google

FRINGED LOOSESTRIFE:

Steironèma ciliàtum.

Primrose Family. June—Aug.

Found in low ground, thickets. Maine to British Columbia, south to Gulf States.

FLOWERS—Yellow, divisions 5, a terra-cotta colored ring in the center.

LEAVES— Opposite, a fringe of hairs borne on leaf stems. 18–24 inches high.

NOTES.

Digitized by Google

Fringed Loosestrife.

Digitized by Google

1. AGRIMONY: *Agrimònia gryposépala.*
Rose Family. June—Aug.
Found in roadsides, borders of woods and fields. Canada to North Carolina, west to California.
FLOWERS—Yellow, small, divisions 5, borne on slender spikes terminating branches, 2–4 feet high.
LEAVES— Composed of usually 7 toothed leaflets, with 1 or more pairs of tiny leaflets between; fragrant when crushed.

2. CELANDINE: Greater Celandine: Swallow-wort. *Chelidònium màjus.*
Poppy Family. May—Aug.
A plant 1–2 feet high, found in roadsides, waste places. Eastern United States. Naturalized from Europe.
FLOWERS—Yellow, on slender stalks, divisions 4. Seed pods erect, 1–2 inches long.
LEAVES— Ornamentally cut and lobed. Stem contains a strong, yellow juice.

NOTES.

Digitized by Google

1. Agrimony. **2. Celandine.**

Digitized by Google

1. LOOSESTRIFE: *Lysimàchia terréstris.*
Primrose Family. June—Aug

Found in low, wet ground. Newfoundland to Northern Georgia, west to Minnesota, Arkansas and Manitoba.

FLOWERS—Yellow, with reddish dots near the center, borne in elongated cluster terminating stem, 1–2 feet high.

LEAVES— Opposite, tapering at both ends.

2. FOUR-LEAVED LOOSESTRIFE; Whorled Loosestrife: Crosswort. *Lysimàchia quadrifòlia.*
Primrose Family. June—Aug.

A slender plant 1–3 feet tall, found in open woods, moist, sandy soil. New Brunswick to Georgia, west to Illinois and Minnesota.

FLOWERS—Yellow, dotted around the center or faintly streaked with red, borne on slender stems springing from the leaf angles.

LEAVES— In whorls of 3–7, usually 4.

NOTES.

Digitized by Google

1. Loosestrife. **2. Four-leaved Loosestrife.**

Digitized by Google

CONE-FLOWER; PURPLE CONE-FLOWER: Black-eyed Susan: Yellow or Ox-eye Daisy: Nigger-head: Golden Jerusalem. *Rudbéckia hírta.*

Composite Family. June—Sept.

A plant 1–3 feet tall, found in fields and open, sunny places. New England to Gulf States, Ontario and Northwest to Colorado.

FLOWER-HEADS—Rays 10–20, yellow or orange-yellow, surrounding a conical, dark purplish-brown disk.

LEAVES— Nearly or quite toothless, lower leaves broader at the tip, 3-ribbed. Leaves and stems rough-hairy.

TALL CONE-FLOWER: Green-headed Coneflower: Thimble-weed. *R. laciniàta.*

Rays 6–10, yellow, slightly drooping, surrounding a greenish yellow disk. Leaves variously divided, lobed and toothed. A plant 3–10 feet tall. Found in moist thickets. Quebec to Montana, south to Gulf States.

NOTES.

Digitized by Google

Cone-flower.

Digitized by Google

1. COMMON MULLEIN: Great Mullein: Velvet or Flannel Plant: Mullein Dock: Aaron's Rod. *Verbáscum Thápsus.*

Figwort Family. June—Sept.

A plant 2–7 feet tall, found in dry fields, waste places. Nova Scotia to Florida, west to Minnesota and Kansas. Naturalized from Europe.

FLOWERS—Yellow, 5-parted, in dense, elongated spike.

LEAVES— Thick, velvety, in a rosette on the ground, others clasping the plant stem.

2. MOTH MULLEIN: *Verbáscum Blattària.*

Figwort Family. June—Sept.

A plant 2–5 feet tall, found in dry fields, waste places. Fairly common through United States and Canada. Naturalized from Europe.

FLOWERS—Yellow, sometimes white, tinged on the back with lavender, fringed in the center with ruddy hairs; borne in loose spike.

LEAVES— Smooth, more or less deeply toothed.

NOTES.

Digitized by Google

1. Common Mullein.

2. Moth Mullein.

Digitized by Google

WILD LETTUCE: Tall Lettuce: Wild Opium: Horse-weed. *Lactùca canadénsis.*

Composite Family. June—Sept.

Found in roadsides, borders of fields, thickets, moist soil. Canada to Georgia, west to Arkansas.

FLOWER-HEADS—Pale-yellow, numerous, small, in loose, spreading cluster terminating stem, 4–10 feet high.

LEAVES— Lower leaves deeply toothed, often 1 foot long, narrowed toward the base; upper leaves lance-shaped, toothless.

NOTES.

Digitized by Google

Wild Lettuce.

Digitized by Google

1. YELLOW or HOP CLOVER:

Trifòlium agràrium.

Pulse Family. June—Sept.

Found in sandy fields, roadsides. Naturalized from Europe. Virginia to Iowa, far northward.

FLOWERS—In heads composed of many tiny, yellow florets, resembling small, dried hops when withered.

LEAVES— In groups of 3 leaflets, almost imperceptibly toothed.

2. LADY'S SORREL: Yellow Wood-sorrel.

Oxalis corniculàta.

Wood Sorrel Family. May—Sept.

Common weed in all situations. Nova Scotia to Gulf, west to Dakota.

FLOWERS—Yellow, small, 5 divisions.

LEAVES— Composed of 3 heart-shaped leaflets.

NOTES.

Digitized by Google

1. Yellow or Hop Clover. **2. Lady's Sorrel.**

Digitized by Google

BLACK MUSTARD: *Brássica nìgra.*
Mustard Family. June—Sept.

A plant 2–5 feet high, found in fields and roadsides. Naturalized from Europe. Common throughout United States.

FLOWERS—Yellow, 4-parted, in loose clusters terminating branches. Seed pods about ½ inch long, 4-sided, lying close to the stem.

LEAVES— Variously lobed and toothed, terminal lobe largest.

YELLOW ROCKET; WINTER CRESS: Herb of St. Barbara: Yellow Bitter, or Rocket Cress. *Barbarèa vulgàris.*
Mustard Family. Apr.—June.

Flowers similar to those of preceding; seed pods long, curved. Upper leaves stemless, lower leaves usually in 5 divisions, terminal one largest. Lake Superior northward and westward; apparently introduced eastward.

NOTES.

Digitized by Google

Black Mustard.

Digitized by Google

Digitized by Google

NOTES.

Digitized by Google

1. COMMON EVENING PRIMROSE: Evening Primrose: Night Willow-herb.

Oenothèra biénnis.

Evening Primrose Family. June—Sept.

A plant 1–5 feet high, found in fields and roadsides. Labrador to Gulf, west to Rockies.

FLOWERS—Yellow, 4-parted, 1–2 inches across, in terminal spikes, opening in the evening.

LEAVES— Alternate, lance-shaped, mostly seated on stem, obscurely toothed. Plant stem leafy, hairy, more or less branched.

2. SUNDROPS: Common Sundrops:

Oenothèra fruticòsa.

Evening Primrose Family. May—July.

Plant 1–3 feet tall, found in fields, roadsides. New England to South Carolina, west to Indiana and Michigan.

FLOWERS—Yellow, ½–1 inch across, 4 divisions, terminating stem and from the leaf angles. Seed pods ribbed and winged.

LEAVES— Oblong to lance-shaped, minutely toothed.

O. pùmila is a similar but smaller species. Leaves toothless, basal leaves broader at the tip. More western in its range.

NOTES.

Digitized by Google

1. Common Evening Primrose. 2. Sundrops.

Digitized by Google

SPOTTED TOUCH-ME-NOT; JEWEL-WEED: Silver Cap: Lady's Ear-drop: Snap Weed: Wild Balsam: Wild Lady's Slipper. *Impàtiens biflòra.*

Touch-me-not Family. June—Sept.

A smooth-stemmed, branching plant, 2–5 feet high, found on brooksides and in moist, shady places. Nova Scotia to Oregon, south to Missouri and Florida.

FLOWERS—Yellow, more or less spotted with reddish brown; sac longer than it is broad, tapering to an incurved spur. Seed pods burst when touched.

LEAVES— Long-stemmed, coarsely toothed.

PALE TOUCH-ME-NOT: *I. pállida.*

Flowers pale yellow, sparingly dotted with reddish brown; sac broader than long with short spur. Most abundant northward.

NOTES.

Digitized by Google

Spotted Touch-me-not.

Digitized by Google

COMMON ST. JOHN'S WORT:

Hypericum perforàtum.

St. John's-wort Family. June—Sept.

A plant 1–2 feet high, found in fields, road-sides. Common throughout the United States. Naturalized from Europe.

FLOWERS—Yellow, in clusters terminating the branches; divisions 5, black-dotted along the margins.

LEAVES— Small, opposite, oblong, more or less black-dotted.

NOTES.

Digitized by Google

Common St. John's-wort.

Digitized by Google

BUTTER-AND-EGGS: Yellow Toad Flax: Flax-weed: Brideweed: Eggs and Bacon.

Linària vulgàris.

Figwort Family. **June—Oct.**

A plant 1–3 feet tall, found in fields, roadsides, waste places. Nova Scotia to Virginia, west to Manitoba and Nebraska. Naturalized from Europe.

Flowers—Yellow and orange, 2-lipped, with a long spur, borne on terminal spikes.

Leaves— Long, narrow, numerous.

NOTES.

Digitized by Google

Butter-and-eggs.

Digitized by Google

1. YELLOW FRINGED ORCHIS:

Habenària ciliàris.

Orchis Family. July—Aug.

Found in moist meadows, sandy bogs. Vermont and Massachusetts to Michigan, Missouri, and southward.

FLOWERS—Golden or orange-yellow, lower lip fringed, borne in spike terminating a stem, 1–2 feet high.

LEAVES— Lance-shaped, clasping.

2. DOWNY FALSE FOXGLOVE:

Gerárdia flàva.

Figwort Family. July—Aug.

A plant 2–4 feet tall, found in open woods. New England to Gulf States, west to Ontario and Wisconsin, Iowa, Arkansas.

FLOWERS—Yellow, trumpet-shaped, 5-lobed.

LEAVES— Opposite, lower ones more or less irregularly lobed and toothed. Plant stem and leaves downy.

SMOOTH FALSE FOXGLOVE: *G. virgínica.*

Flowers slightly larger than those of preceding. Lower leaves deeply cut and toothed. Whole plant smooth.

NOTES.

Digitized by Google

1. Yellow Fringed Orchis. 2. Downy False Foxglove.

Digitized by Google

ELECAMPANE: Horseheal: Yellow Starwort.
Inula Helènium.

Composite Family. July—Sept.

Plant 2–6 feet tall. Found in roadsides, fields, damp pastures. Naturalized from Europe. Minnesota and Missouri east to Nova Scotia, south to South Carolina.

FLOWER-HEADS—Yellow, solitary or few, 2–4 inches across, rays long and narrow.

LEAVES— Large, white-veined, rough above, woolly beneath, toothed; lower leaves stemmed, upper ones partly clasping plant stem.

NOTES.

Digitized by Google

Elecampane.

Digitized by Google

1. PARTRIDGE PEA: Large-flowered Sensitive Plant: *Cássia Chamaecrísta.*
Pulse Family. July—Sept.
A plant 1–2 feet tall, found in dry or sandy fields. Massachusetts to Minnesota and southward.
FLOWERS—Yellow, 5-parted, about 1 inch across, 2–4 from the bases of the leaves.
LEAVES— Composed of 10–15 pairs of leaflets.

WILD SENNA: *C. marilándica.*
Flowers yellow, divisions 5, 3 forming an upper lip, 2 a lower one. Leaflets 5–9 pairs. 3–8 feet high. Moist soil, swamps. New England to Ohio, Tennessee, and North Carolina.

2. FALL DANDELION: Autumnal Hawkbit. *Leóntodon autumnàlis.*
Composite Family. July—Nov.
Found in fields, roadsides. Throughout northern United States, south to Ohio and New Jersey and Pennsylvania. Naturalized from Europe.
FLOWER-HEADS—Yellow, smaller than the common dandelion, on slender, branching stalks, 6–18 inches high.
LEAVES— Long, narrow, more or less coarsely toothed.

Digitized by Google

1. Partridge Pea. **2. Fall Dandelion.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

TANSY: Bitter-buttons. ***Tanacètum vulgàre.***
Composite Family. **July—Sept.**

Found in roadsides, waste places and around old buildings. Introduced from Europe. Nova Scotia to North Carolina and Missouri, west to Minnesota.

Flower-heads—Yellow, in rather flat clusters terminating stem, $1\frac{1}{2}$–3 feet tall.

Leaves— Deeply cut into narrow, toothed divisions, aromatic.

NOTES.

Digitized by Google



Tansy.

Digitized by Google

ORANGE or TAWNY HAWKWEED; DEVIL'S PAINT-BRUSH: Grim the Collier.

Hieràcium aurantìacum.

Composite Family. July—Sept.

Found in fields, pastures. Naturalized from Europe. Canada south to Pennsylvania and Middle Atlantic States.

FLOWER-HEADS—Reddish orange, in a group terminating stem, 6–20 inches high.

LEAVES— Mostly at the base, blunt, lance-shaped. Whole plant hairy.

CANADA HAWKWEED: *H. canadénse.*

FLOWERS—Yellow, terminating stem 1–4 feet tall.

LEAVES— Seated on stem, coarsely toothed, especially the lower ones.

NOTES.

Digitized by Google



Orange Hawkweed.

Digitized by Google

1. HAWKWEED: Rough Hawkweed.
Hieràcium scàbrum.

Composite Family. July—Sept.

A plant 1–3 feet high, found in dry woods, pastures. Nova Scotia to Georgia, west to Nebraska.

FLOWER-HEADS—Yellow, in branching, terminal clusters.

LEAVES— Blunt, almost toothless, seated on stem. Whole plant hairy.

HAIRY HAWKWEED: *H. Gronòvii.*

A similar plant with slender stem, very leafy and hairy below.

2. RATTLESNAKE-WEED: Early or Vein-leaf Hawkweed: Snake or Poor Robin's Plantain. *Hieràcium venòsum.*

Composite Family. June—Sept.

Found in dry woods, open, sandy places. Atlantic States to Minnesota and Iowa.

FLOWER-HEADS—Yellow, resembling small dandelions, few on slender, branching stalk, 10–20 inches high.

LEAVES— In a rosette, hairy, with purplish veins.

NOTES.

Digitized by Google

1. Hawkweed. **2. Rattlesnake-weed.**

Digitized by Google

1. THIN-LEAVED or TEN-PETALED SUNFLOWER: *Helianthus decapetalus.*

Composite Family. Aug.—Sept.

A plant 2–5 feet tall, found in moist woods, beside streams. Eastern United States to Minnesota, Missouri and southward.

FLOWER-HEADS—Yellow, 2–3 inches broad, rays about 10.

LEAVES— Egg-shaped, pointed, opposite, smooth or slightly rough. Plant stem rough above; smooth below, 2–5 feet tall.

2. LARGER BUR-MARIGOLD: Brook Sunflower. *Bidens laevis.*

Composite Family. Aug.—Oct.

Plant 1–2 feet high, found in swamps, wet places. New England to Minnesota and southward.

FLOWER-HEADS—Yellow, 1–2½ inches across, rays 8–10. Seed vessels with 2–4 prongs.

LEAVES— Opposite, toothed, narrow lance-shaped, seated on stem.

NOTES.

Digitized by Google

1. Thin-leaved Sunflower. **2. Larger Bur-Marigold.**

Digitized by Google

NOTES.

Digitized by Google

GOLDEN CORYDALIS: *Corýdalis aúrea.*
Fumitory Family. Mch.—May.

A smooth-stemmed, branching plant 6–14 inches high, found in woodlands. Nova Scotia to Pennsylvania, west to the Rocky Mountains.

FLOWERS—Golden yellow, irregular, about ½ inch long, borne in short, terminal clusters. Seed pods drooping or spreading, slightly curved, lumpy.

LEAVES— Compound, finely dissected.

ROUND-LEAVED, or EARLY YELLOW VIOLET: *Viola rotundifòlia.*
Violet Family. Apr.—May.

Found in woodlands. A plant 2–4 inches high. New England to Georgia.

FLOWERS—Yellow, with purplish veins, small, side divisions bearded.

LEAVES— Round or long heart-shaped, indistinctly toothed.

Digitized by Google

DOWNY YELLOW VIOLET: *Viola pubéscens.*

Violet Family. Apr.—May.

An erect leafy plant 6–15 inches high, common in dry woodlands. Maine to Ontario, south to Maryland and Kansas.

FLOWERS—Golden yellow, with purplish veins, borne on stalks from the leaf angles.

LEAVES— Broadly heart-shaped, somewhat toothed. Plant stem and leaves slightly downy.

GOLDEN CLUB: *Orónti um aquáticum.*

Arum Family. Apr.—May.

Found in shallow water, swamps, chiefly near the Atlantic coast. New England to Gulf of Mexico.

FLOWERS—Yellow, minute, crowded on a club 1–2 inches long, terminating stalk 6–36 inches high; the stalk flattened just below the flower cluster.

LEAVES— Dark green, oblong, long-stemmed, floating or erect.

Digitized by Google

INDIAN POKE; AMERICAN WHITE HELLEBORE: Itch-weed. *Veràtrum víride.*

Lily Family. Apr.—June.

A plant 2–7 feet high, poisonous throughout; found in low, wet places. Canada to British Columbia, south to Georgia, Tennessee, Minnesota.

FLOWERS—Very small, forming a large, terminal, yellow-green cluster.

LEAVES— Large, broad, with an uneven surface, soon turning brown and unsightly.

YELLOW MELILOT or SWEET CLOVER: *Melilòtus officinàlis.*

Pulse Family. June—Aug.

Found in waste places. Naturalized from Europe.

FLOWERS—Resemble the White Melilot, except in color, yellow, small, in slender spikes terminating branches, 2–4 feet high.

LEAVES— Small, in groups of 3, blunt tipped, toothed.

Digitized by Google

FROSTWEED; ROCKROSE: Frost-flower, Frost-wort: Long-branched Frost-weed: Canadian Rockrose. ***Heliánthemum canadénse.***

Rockrose Family. **June—Aug.**

Found in sandy soil. New England to South Carolina, west to Wisconsin.

Flowers—Of 2 kinds: the earlier ones solitary, about 1 inch across, yellow, 5 divisions; later ones small, clustered at the base of the leaves.

Leaves— Lance-oblong, pale beneath, set close to the stem.

NOTES.

Digitized by Google

BUTTERFLY WEED; PLEURISY ROOT: Orange-root: Orange Milkweed.

Asclèpias tuberòsa.

Milkweed Family. June—Sept.

Found in dry fields. Maine to Gulf States, west to Ontario and Arizona. Common southward.

FLOWERS—Similar in structure to the Common Milkweed, reddish orange or orange-yellow, in erect clusters terminating stem, 2–3 feet high.

LEAVES— Narrow-oblong, or lance-shaped, toothless, hairy beneath, nearly or quite stemless.

GINSENG: *Pànax quinquefòlium.*

Ginseng Family. July—Aug.

Found in rich woods. Quebec to Alabama, west to Nebraska.

FLOWERS—Inconspicuous, greenish yellow, in a cluster terminating stem, 8–15 inches high. Berries red in a cluster.

LEAVES— Consisting of 3 sets in a whorl, each set composed of 5 somewhat egg-shaped, sharp-pointed, toothed leaflets.

Digitized by Google

HORSE BALM; RICH-WEED: Horse-weed: Stone-root: Citronella.

Collinsònia canadénsis.

Mint Family. July—Sept.

A plant 2–4 feet tall, found in rich, moist woods. New England to Florida, Ontario and Wisconsin to Kansas and Missouri.

FLOWERS—Pale yellow, about ½ inch long, with 3 long, protruding filaments, lemon-scented; 5-lobed, one lobe fringed and much larger than the others; borne in loose, spreading cluster.

LEAVES— Oval, toothed. Plant stem branching.

SNEEZEWEED; SWAMP SUNFLOWER:

Helènium autumnàle.

Composite Family. Aug.—Sept.

A plant 2–6 feet high, found in wet meadows, river banks. Quebec to Florida, west to Arizona, Minnesota and northwest.

FLOWER-HEADS—Yellow, 1–2 inches across, rays drooping, broad, 3–5 cleft, in terminal clusters.

LEAVES— Lance-shaped or oblong, alternate, toothed.

Digitized by Google

FERN-LEAVED FALSE FOXGLOVE: *Gerárdia pediculària.*

Figwort Family. Aug.—Sept.

A plant 1–3 feet high, found in borders of dry woods, thickets. Maine to Ontario, Minnesota, and West Virginia.

FLOWERS—Yellow, bell-shaped, with 5 spreading, rounded lobes, 1 inch or more across, somewhat finely hairy.

LEAVES— Fern-like, deeply cut into many toothed lobes. Plant stem leafy, branching, somewhat sticky-hairy.

GOLDEN ASTER: *Chrysópsis mariàna.*

Composite Family. Aug.—Oct.

Found in dry, sandy soil. Gulf of Mexico, north to Pennsylvania and Long Island.

FLOWER-HEADS—Yellow, aster-like, nearly 1 inch across, in loose, spreading clusters.

LEAVES— Oblong, seated on stem, toothless. Leaves and plant stem silky-hairy when young, becoming smooth with age.

GOLDEN-ROD. The golden-rods are familiar to everyone. They comprise a very extensive group, with many intergrading forms, and none but students care to puzzle out the different species.

Digitized by Google

NOTES.

Digitized by Google

III

RED, PINK, CRIMSON AND MAGENTA

Digitized by Google

SPRING BEAUTY: Claytonia.

Claytònia virgínica.

Purslane Family. Mch.—May.

A plant 6–12 inches high, found in moist, open woods, low meadows. Common west and south.

FLOWERS—White veined with pink, or pink with veins of deeper pink; divisions 5.

LEAVES— Long and narrow, width variable. Roots tuberous.

C. caroliniàna is similar to the preceding, except that the flowers are smaller and fewer, and leaves more oblong.

Digitized by Google

Spring Beauty.

Digitized by Google

1. SHOOTING STAR; AMERICAN COWSLIP: Pride of Ohio. *Dodecàtheon Meàdia.*

Primrose Family. Apr.—June.

Found in open woods, prairies, moist cliffs. Wisconsin to Pennsylvania, south to Texas and Georgia.

FLOWERS—Magenta-pink or white, the 5 divisions turned backward, pendulous, grouped on stalk 8–20 inches high.

LEAVES— Oblong, narrowed toward the base.

2. ARETHUSA: Indian Pink. *Arethùsa bulbòsa.*

Orchis Family. May—June.

Found in bogs and swamps. Newfoundland south to South Carolina mountains, west to Indiana and Minnesota.

FLOWERS—Magenta-crimson, lower lip spotted with magenta crimson and crested with 3 white or yellow hairy ridges; solitary, on stalk 5–10 inches high.

LEAVES— Long and narrow, solitary, appearing after the flower opens.

Root bulbous.

Digitized by Google

1. Shooting Star. 2. Arethusa.

Digitized by Google

WAKE ROBIN: Purple Trillium: Ill-scented Wake Robin: Birthroot. *Trillium eréctum.*
Lily Family. Apr.—June.
Plant 8–16 inches high, found in rich woods. Eastern Quebec south to North Carolina, west to Minnesota and Missouri.
FLOWERS—Brown-purple, dull pink, white or greenish, divisions 3, on short stalk from junction of leaves.
LEAVES— Broad, 3 in a whorl.

SESSILE-FLOWERED WAKE ROBIN: Stemless Trillium. *T. séssile.*
Flowers stemless, dark purplish or greenish.
T. recurvàtum. Outer divisions of flower reflexed; colored divisions narrowed at the base; leaves narrowed into a stem.

NOTES.

Digitized by Google

Wake Robin.

Digitized by Google

WILD COLUMBINE: Honeysuckle.

Aquilègia canadénsis.

Crowfoot Family. Apr.—June.

A slender plant 1–2 feet high, found in open, rocky woodlands, pastures. Common throughout the United States.

FLOWERS—Red outside, yellow within; 5 tubes terminating in spurs; borne on slender stalks from the leaf angles.

LEAVES— Compound, more or less divided, teeth rounded.

NOTES.

Digitized by Google

Wild Columbine

Digitized by Google

1. WILD PINK: Catchfly. *Silène pennsylvánica.*
Pink Family. Apr.—June.
Found in gravelly, rocky places. Eastern New England to New York and Kentucky, south to Georgia.
FLOWERS—Crimson-pink, divisions 5, somewhat sticky immediately below the flower, in terminal clusters, 4–10 inches high.
LEAVES— Blunt lance-shaped, mostly at the base; a few opposite, lance-shaped leaves seated on the stems.

2. MAIDEN PINK: *Diánthus deltoìdes.*
Pink Family. June—Aug.
Found in fields, waste places. New England to Michigan. Naturalized from Europe.
FLOWERS—Crimson-pink or white, 5 toothed divisions, blooming singly, terminating slender stems 6–12 inches high.
LEAVES— Narrow, erect.

DEPTFORD PINK: *D. Armèria.*
Flowers crimson-pink, white-dotted, divisions toothed. Plant stem and leaves finely hairy. Naturalized from Europe.

Digitized by Google

1. Wild Pink. **2. Maiden Pink.**

Digitized by Google

1. WILD CRANESBILL; WILD or SPOTTED GERANIUM:

Geránium maculátum.

Geranium Family. Apr.—July.

A plant 1–2 feet high, found in open woods, roadsides. Maine to Manitoba and southward.

FLOWERS—Pale or deep magenta-pink or lavender, divisions 5.

LEAVES— Usually 5-parted variously toothed, stem leaves opposite. Leaves and plant stem hairy.

2. HERB ROBERT: Red Robin: Red Shanks: Dragon's Blood. *Geránium Robertiánum.*

Geranium Family. June—Oct.

Found in moist, rocky woods. Eastern Quebec to Minnesota, south to New Jersey and Missouri.

FLOWERS—Deep or pale magenta or pink, about ½ inch across, divisions 5.

LEAVES— 3–5 parted, variously cleft and toothed, strongly scented.

Stems spreading, hairy, 6–18 inches high.

Digitized by Google

1. Wild Cranesbill. **2. Herb Robert.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

PITCHER-PLANT: Indian Dipper: Side-saddle Flower: Huntsman's Cup.

Sarracènia purpùrea.

Pitcher-plant Family. May—June.

Found in mossy bogs. Labrador to Minnesota, Rocky Mountains, southeast of Alleghenies to Florida.

FLOWERS—Variable in color, the 5 outer divisions greenish to madder purple, inner divisions dull pink; solitary, nodding, borne on stalk 1–2 feet high.

LEAVES— Hollow, usually green with reddish purple veins.

NOTES.

Digitized by Google



Pitcher-plant.

Digitized by Google

STEMLESS, PINK, or VENUS' LADY'S SLIPPER: Moccasin Flower. *Cyripèdium acaúle.*
Orchis Family. May—June.
Found in dry woods. Newfoundland to North Carolina, west to Indiana, Michigan, Minnesota.
FLOWERS—Pouch variable crimson, veined with deeper crimson, sometimes white, solitary, on stalk 6–12 inches high.
LEAVES— 2, from the base, 6–8 inches long.

NOTES.

Digitized by Google



Stemless Lady's Slipper.

Digitized by Google

SHOWY ORCHIS: Gay, or Spring Orchis.

Orchis spectábilis.

Orchis Family. **May—June.**

Found in rich woods. New Brunswick to Georgia, west to Minnesota and Missouri.

Flowers—Magenta, lower lip almost white, 3–6 borne on stalk 4–12 inches high.

Leaves— 2, broad, shining, from the base.

NOTES.

Digitized by Google

Showy Orchis.

Digitized by Google

WILD AZALEA: Pinxter Flower: Wild Honeysuckle: Pink or Purple Azalea.

Rhododéndron nudiflòrum.

Heath Family. **May—June.**

A shrub 2–6 feet high, found in open woods, swamps. Massachusetts to Florida, west to Illinois, Missouri, Texas.

Flowers—Crimson-pink to nearly white, the base of the long, narrow tube deeper in color, divisions 5, 6 long, protruding filaments; flowers borne in small, terminal clusters, opening with or before the leaves.

Leaves— Usually clustered, tapering at each end.

NOTES.

Digitized by Google

Wild Azalea

Digitized by Google

1. FRINGED MILKWORT; FLOWERING WINTERGREEN: Gay Wings: Fringed Polygala. *Polýgala paucifòlia.*

Milkwort Family. May—July.

Found in moist woods. Eastern Quebec to Minnesota and Illinois, south in Alleghenies.

FLOWERS—Magenta, magenta-crimson, or rarely white, very irregular in form, fringed on the tip.

LEAVES— Clustered at the summit of the stem, 3–6 inches high.

2. FIELD MILKWORT: Common or Purple Milkwort: Purple Polygala. *Polýgala sanguínea.*

Milkwort Family. June—Sept.

Found in moist or sandy fields, roadsides. Southern Canada to North Carolina, west to the Mississippi.

FLOWERS—Deep or pale magenta, rarely white, tiny, clustered in globular or oblong heads, terminating stems 6–12 inches high.

LEAVES— Small, narrow

Digitized by Google

1. Fringed Milkwort.

2. Field Milkwort.

Digitized by Google

MOUNTAIN or AMERICAN LAUREL: Calico Bush: Spoonwood: Calmoun: Broad-leaved Kalmia. *Kálmia latifòlia.*

Heath Family. May—July.

Found in rocky woods. Canada and Maine mountains to Gulf of Mexico, west to Ohio and Tennessee. Usually a shrub 3–6 feet high, but in its southern range, often tree-like.

FLOWERS—White or pink, bowl-shaped, 5-pointed; 10 filaments spring from the center, their tips lodged in little pockets; flowers borne in clusters terminating the branches.

LEAVES— Long, evergreen, tapering at both ends.

NOTES.

Digitized by Google



Mountain Laurel.

Digitized by Google

TWISTED-STALK: Sessile-leaved Twisted Stalk.
Stréptopus ròseus.

Lily Family. May—July.

A plant 1–2½ feet high, found in moist woods. Newfoundland to Minnesota, south to Ohio, Pennsylvania, mountains of Georgia.

FLOWERS—Bell-shaped, dull, purplish pink, solitary, from the axils of the leaves.

LEAVES— Alternate, seated on stem.

CLASPING-LEAVED TWISTED-STALK:
S. amplexifòlius.

Differs from the preceding in its greenish flowers with widely spreading segments, and whitish bloom on the under side of the leaves. Leaves strongly clasping.

NOTES.

Digitized by Google

Twisted-stalk.

Digitized by Google

1. WOOD BETONY; LOUSEWORT: Beefsteak Plant: High Heal-all. *Pediculàris canadénsis.*
Figwort Family. May—July.
Found in open woods. Nova Scotia to Florida, west to Manitoba, Kansas, Colorado.
FLOWERS—Greenish yellow and purplish red, 2-lipped, in short, dense clusters, terminating stems 6–18 inches high.
LEAVES— Long, narrow, deeply toothed.

2. BOUNCING BET; SOAPWORT: Hedge Pink: Bruisewort: Old Maid's Pink: Fuller's Herb. *Saponària officinàlis.*
Pink Family. June—Sept.
A plant 1–2 feet high, found in roadsides, waste places. Adventive from Europe. Common, escaped from gardens.
FLOWERS—Magenta-pink to white, divisions 5, but often double, borne in terminal clusters.
LEAVES— Opposite, toothless, 3–5-ribbed. Plant stem stout, thick jointed.

Digitized by Google

1. Wood Betony. **2. Bouncing Bet.**

Digitized by Google

PALE CORYDALIS: *Corýdalis sempérvirens.*
Fumitory Family. May—Aug.

Found in rich, rocky places. Eastern Quebec to Alaska, south to Georgia, Kentucky, and Montana.

FLOWERS—Crimson-pink about ½ inch long, in loose clusters terminating branches, 1–2 feet high.

LEAVES— Compound, variously cut and divided.

NOTES.

Digitized by Google

Pale Corydalis.

Digitized by Google

SHEEP LAUREL; LAMBKILL: Wicky: Sheep-poison: Calf-kill: Narrow-leaved Laurel. *Kálmia angustifòlia.*

Heath Family. June—July.

A shrub 1–3 feet high, found on hillsides and in swamps. Labrador and Ontario to Michigan, south to Georgia.

FLOWERS—Crimson-pink, in clusters encircling plant stem.

LEAVES— Narrow, opposite, or in groups of 3. Stem terminated by the upright newer leaves.

PALE LAUREL: Swamp Laurel. *K. polifòlia.*

May be distinguished by its flowers terminating the stems, and by its leaves, which are usually set opposite, edges rolled back, whitish beneath. Found in cold bogs and hillside swamps. Labrador to Alaska, south to Pennsylvania and Michigan, west to Pacific coast.

Digitized by Google

Sheep Laurel.

Digitized by Google

1. GRASS PINK: Calopogon.
Calopògon pulchéllus.
Orchis Family. June—July.
Found in open bogs, meadows. Newfoundland to Florida, west to Minnesota and Missouri.
FLOWERS—Magenta-pink, the lip on the upper side of flower densely bearded with yellow and magenta-crimson hairs; 3–12 flowers terminate stalk 10–16 inches high.
LEAF— Solitary, grass-like, from bulb.

2. ROSE, or SWEET POGONIA; SNAKE MOUTH: *Pogònia ophioglossoìdes.*
Orchis Family. June—July.
Found in wet meadows, swamps. Newfoundland to Florida, west to Indiana and Minnesota.
FLOWERS—Crimson-pink, crest of lower lip yellow to white, solitary, terminating stem 8–15 inches high.
LEAVES— A small leaflet borne just below the flower, and a larger leaf midway of the stem.

Digitized by Google

1. Grass Pink.

2. Rose Pogonia.

Digitized by Google

WILD ORANGE-RED LILY; WOOD LILY; Flame or Philadelphia Lily.

Lílium philadélphicum.

Lily Family. June—July.

Found in dry, sandy soil, woodland borders. New England to North Carolina, west to Minnesota and Missouri.

FLOWERS—Orange or reddish orange, spotted inside with purplish brown; 1–3 borne erect at the summit of stem 1–3 feet high.

LEAVES— In whorls of 3–8.

NOTES.

Digitized by Google

Wild Orange-red Lily.

Digitized by Google

1. SPREADING DOGBANE: Fly-trap Dogbane: Bitter-root: Honey-bloom: Indian Hemp.
Apócynum androsaemifòlium.

Dogbane Family. June—Aug.

Found in open woods, roadsides. Northern Canada to Georgia, west to Nebraska.

FLOWERS—Pink tinted, veined inside with deeper pink, bell-shaped with 5 spreading lobes, in loose clusters on the tips of the branches.

LEAVES— Opposite, toothless, short-stemmed. Plant stem branching, shrubby, 1–4 feet high.

2. TWIN-FLOWER: Ground Vine.
Linnæa boreàlis.

Honeysuckle Family. June—Aug.

A trailing vine, stems 6–20 inches long, found in moist, mossy woods. Labrador to New Jersey and mountains of Maryland, west to Minnesota; also far west.

FLOWERS—Whitish, tinged and striped with crimson-pink, fragrant, slender bell-shaped, 5-lobed, borne in pairs, on stalk 3–4 inches high.

LEAVES— Opposite, rounded, scallop-toothed, evergreen.

Digitized by Google

1. Spreading Dogbane. **2. Twin-flower.**

Digitized by Google

1. MEADOW-SWEET: Queen-of-the-Meadow: Quaker Lady. *Spiraèa latifòlia.*

Rose Family. June—Aug.

A shrubby plant, 2–4 feet high, found in rocky pastures, borders of fields. Newfoundland to Georgia, west to Minnesota and Missouri.

FLOWERS—Small, flesh pink or whitish, in dense, terminal clusters.

LEAVES— Light green, oval or oblong, rather coarsely toothed. Plant stems red or purplish brown.

S. salicifòlia differs from the preceding in its white flowers, yellowish brown stems, and finely toothed leaves. Chiefly in low ground.

2. HARDHACK; STEEPLE BUSH: *Spiraèa tomentòsa.*

Rose Family. July—Sept.

A shrubby plant, with erect, woolly stems 2–4 feet high, found in low ground or on hillsides. New Brunswick to Georgia, west to Minnesota and Kansas.

FLOWERS—Pink or magenta, rarely white, in dense, steeple-like clusters.

LEAVES— Dark green above, whitish-woolly beneath, toothed.

Digitized by Google

1. Meadow-sweet. **2. Hardhack.**

Digitized by Google

PURPLE FLOWERING RASPBERRY: Virginia Raspberry. ***Rùbus odoràtus.***

Rose Family. **June—Aug.**

A sprawling shrub, 3–5 feet high, found in rocky woods, shaded roadsides. Nova Scotia to Georgia, west to Michigan.

FLOWERS—Crimson-pink, fading as they get older, 1–2 inches broad. Fruit a thin, red, scarcely edible berry.

LEAVES— Large, lobed and toothed, hairy, similar to those of the maple in shape. Plant stems hairy.

NOTES.

Digitized by Google

Purple Flowering Raspberry.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

1. LARGE, or EARLY, PURPLE FRINGED ORCHIS: *Habenària fimbriàta.*

Orchis Family. June—Aug.

Found in wet woods, swamps. Newfoundland to North Carolina, west to Michigan.

FLOWERS—Magenta-pink, varying to almost white, lower lip 3-parted, deeply fringed, in spike terminating stem 1–5 feet high.

LEAVES— Lance-shaped above, somewhat oval below.

SMALLER PURPLE FRINGED ORCHIS: *H. psycòdes.*

Bears smaller blossoms, less deeply fringed. A plant 1–3 feet high, found in similar situations west to Indiana and Minnesota.

2. PRINCE'S PINE; PIPSISSEWA: *Chimáphila umbellàta.*

Heath Family. June—Aug.

Found in dry woods. Nova Scotia to Georgia, west to Pacific.

FLOWERS—Flesh color or pinkish, waxy, 5-parted, in small groups terminating stem 3–10 inches high.

LEAVES— Dark, shining green, evergreen, sharply toothed, opposite or in whorls.

SPOTTED WINTERGREEN: Pipsissewa. *C. maculàta.*

Leaves white veined, tapering toward the tip. Found in similar situations west to Minnesota and Mississippi.

Digitized by Google

1. Purple Fringed Orchis. **2. Prince's Pine.**

Digitized by Google

COMMON MILKWEED: Silkweed.
Asclèpias syrìaca.

Milkweed Family. June—Aug.

A plant 3–5 feet high, found in fields, roadsides, waste places. Common. New Brunswick south to North Carolina and Kansas; also westward.

FLOWERS—Variable in color, dull crimson-pink, lilac or yellowish; one or more thick clusters springing from the leaf angles. Seed pods large, filled with silky down and flat, overlapping seeds.

LEAVES— Opposite, broad oblong, toothless.

PURPLE MILKWEED: *A. purpurascens.*

Similar in habit to the preceding. Found in woods, dry fields. New Hampshire to Ontario, Minnesota, Kansas, and southward.

FLOWERS—Crimson or crimson-magenta, in terminal clusters.

LEAVES— Elliptical or egg-shaped, upper ones taper-pointed, minutely downy beneath.

NOTES.

Digitized by Google

Common Milkweed.

Digitized by Google

HEDGE or GREAT BINDWEED; WILD MORNING-GLORY: Bell-bind: Rutland Beauty: Lady's Nightcap. *Convólvulus sèpium.*

Convolvulus Family. June—Sept.

A vine 3–10 feet long, found in thickets, fields, waysides. North Atlantic States westward.

FLOWERS—Pink or white, large, bell-shaped, on long stems from leaf angles.

LEAVES— Triangular or arrowhead-shaped, long-stemmed.

FIELD BINDWEED: *C. arvénsis.*

Similar to the preceding, but much smaller. Flowers not over 1 inch long, pinkish or white, often in pairs. Naturalized from Europe. North Atlantic States west to Kansas.

NOTES.

Digitized by Google



Hedge Bindweed.

Digitized by Google

1. LARGE MARSH PINK:
Sabàtia dodecándra.
Gentian Family. July—Aug.
A plant 1–2 feet high, found on borders of brackish ponds near the Atlantic coast. Massachusetts to North Carolina.
FLOWERS—Crimson-pink, about 2 inches across, 8–12 divisions.
LEAVES— Lower leaves blunt-tipped, tapering toward the base, upper leaves narrow, lance-shaped.

BITTER BLOOM: Rose Pink: Square-stemmed Sabatia: Rose Centaury. *S. angulàris.*
Flowers crimson-pink or white, greenish star in the center, 5-parted. Leaves opposite, 5-veined, clasping the square plant stem. Found in rich soil. New York to Florida, west to Indian Territory, Michigan, Ontario.

2. SEA or MARSH PINK: Rose of Plymouth.
Sabàtia stellàris.
Gentian Family. July—Sept.
Found along the Atlantic coast.
FLOWERS—Crimson-pink, about 1 inch across.
LEAVES— Upper leaves narrow, widening as they approach the root.

SLENDER MARSH PINK: *S. grácilis.*
Resembles the Sea Pink, but stems are more slender, leaves narrower.

Digitized by Google



1. Large Marsh Pink.



2. Sea Pink.

Digitized by Google

SWAMP MILKWEED: *Asclèpias incarnàta.*
Milkweed Family. July—Aug.

Found in swampy places. New Brunswick, west and southwest.

FLOWERS—Dull crimson or crimson-pink, in clusters, terminating stem 2–4 feet high.

LEAVES— Narrow, lance-shaped, short-stemmed; 2 downy lines on stems below flower clusters.

FOUR-LEAVED MILKWEED:
A. quadrifòlia.

Bears 1–2 whorls of 4 leaves each near the middle of plant stem. Flowers pinkish and white. New Hampshire to North Carolina, Ontario to Arkansas.

NOTES.

Digitized by Google

Swamp Milkweed.

Digitized by Google

1. GREAT or SPIKED WILLOW-HERB; FIREWEED: *Epilòbium angustifòlium.*
Evening Primrose Family. July—Aug.
Found in dry soil, clearings, newly burnt land. New England to North Carolina, west to Minnesota and Kansas, and, with few exceptions, to the Pacific.
FLOWERS—Magenta, rarely white, divisions 4, in spike terminating stem 2–8 feet high.
LEAVES— Narrow, quite toothless, tapering.

2. CARDINAL FLOWER: *Lobèlia cardinàlis.*
Lobelia Family. Aug.—Sept.
Found in low, moist ground. New Brunswick to Ontario and southward.
FLOWERS—Rich, deep red, lower lip 3-lobed, in somewhat 1-sided spike terminating stem 2–4 feet high.
LEAVES— Dark green, oblong to lance-shaped, slightly toothed, upper leaves stemless.

Digitized by Google

1. Great Willow-herb. 2. Cardinal Flower.

Digitized by Google

1. MEADOW BEAUTY; DEER-GRASS:

Rhéxia virgínica.

Meadow Beauty Family. July—Sept.

A square-stemmed plant 10–18 inches high, found in sandy marshes or near water. Maine to Florida, Ontario to Iowa and southward.

FLOWERS—Magenta or purplish, division 4, 8 prominent, golden yellow, floral organs.

LEAVES— Opposite, mostly 5-veined, toothed, seated on stem.

MARYLAND MEADOW BEAUTY:

R. mariàna.

May be distinguished from the preceding by its round stems and short-stemmed leaves. Found south of New Jersey.

2. OSWEGO TEA; BEE BALM:

Monárda dídyma.

Mint Family. July—Sept.

A square-stemmed plant 2–3 feet high, found in moist ground, especially near streams, from New England west and south; also cultivated in gardens.

FLOWERS—Scarlet, 2-lipped, clustered in terminal heads, with a number of leaflets below.

LEAVES— Opposite, toothed, dark green, aromatic. Plant stem hairy.

Digitized by Google

1. Meadow Beauty. **2. Oswego Tea.**

Digitized by Google

1. PINK KNOTWEED; SMARTWEED: Common Persicaria: Jointweed.

Polýgonum pennsylvánicum.

Buckwheat Family. July—Sept.

A plant of sprawling, branching habit, 1–3 feet high, common in waste places, moist soil.

FLOWERS—Pink or greenish, small, crowded on short, narrow, terminal spikes.

LEAVES— Long, narrow, toothless, tapering toward the tip. Plant stems enlarged at the joints.

LADY'S THUMB: *P. Persicària.*

Is a similar species, leaves usually marked with a dark triangle. Found throughout North America. Naturalized from Europe.

2. WATER PERSICARIA:

Polýgonum amphíbium.

Buckwheat Family. July—Aug.

Found in shallow water. Quebec to New Jersey, west to Pacific.

FLOWERS—Pink, small, in short, dense, terminal clusters.

LEAVES— Usually floating, smooth, shining above, elliptical or oblong.

Digitized by Google

1. Pink Knotweed. **2. Water Persicaria.**

Digitized by Google

JOE-PYE WEED: Trumpet Weed: Gravel or Kidney-root: Purple Thoroughwort: Tall or Purple Boneset. *Eupatòrium purpùreum.*

Composite Family. Aug.—Sept.

A tall plant 3–10 feet high, common in low, swampy places. New England to the Gulf of Mexico and westward.

FLOWER-HEADS—Dull magenta or magenta-crimson, in broad, terminal cluster.

LEAVES— Lance-shaped, toothed, in whorls of 3–6. Plant stem green or purplish.

NOTES.

Digitized by Google

Joe-Pye Weed.

Digitized by Google

RABBIT'S-FOOT, OLD-FIELD, PUSSY, or STONE CLOVER: *Trifòlium arvénse.*
Pulse Family. **Aug.—Sept.**

A plant 4–10 inches high, with branching stem, found in fields, waste places, dry soil. Naturalized from Europe.

Flowers—In soft, fuzzy heads, gray or gray-pink in effect.

Leaves— In groups of 3 leaflets, widest toward the tip.

NOTES.

Digitized by Google

Rabbit's-foot Clover.

Digitized by Google

LIVE-FOR-EVER; GARDEN ORPINE: Midsummer-men: Pudding-bag Plant: Live-long: Witches' Money: Garden-stonecrop.

Sèdum purpùreum.

Orpine Family. **Aug.—Sept.**

Found in fields, waysides. Quebec to Maryland, Michigan and westward. Introduced from Europe.

FLOWERS—Small, dull garnet red or reddish purple, in thick, terminal clusters.

LEAVES— Thick, smooth, toothed, seated on stem 10–20 inches high.

NOTES

Digitized by Google

Live-for-ever.

Digitized by Google

Digitized by Google

NOTES.

Digitized by Google

GROUND or MOSS PINK: *Phlóx subulàta.*
Polemonium Family. Apr.—June.

A low plant, forming moss-like mats, sometimes carpeting the ground. Found in rocky ground, hillsides. Southern New York to Michigan, south to Florida and Kentucky.

FLOWERS—Crimson-pink, magenta to white, 5-parted, divisions usually notched on the tip, borne in small clusters terminating the stems.

LEAVES— Small, narrow, about ½ inch long, growing in tufts at the joints of the plant stems, which are 2–5 inches long.

DOWNY PHLOX: *Phlóx pilòsa.*
Polemonium Family. May—June.

A downy-stemmed plant 1–2 feet high, found in dry, sandy soil. Ontario to Manitoba, New Jersey, Florida, Arkansas and Texas.

FLOWERS—Borne in terminal clusters, crimson-pink, purplish or white, 5-parted; somewhat sticky immediately beneath the flower.

LEAVES— Narrow or lance-shaped, toothless, stemless.

Digitized by Google

VIOLET WOOD SORREL: *Oxalis violàcea.*
Wood Sorrel Family. May—June.

Found in rocky places, open woods. Massachusetts to Minnesota and southward.

FLOWERS—Pale magenta, sometimes white, 5 divisions, borne in clusters of 3–12 terminating a slender stalk 4–8 inches high.

LEAVES— Composed of 3 clover-like leaflets borne on slender stems springing from the root.

HAIRY BEARD-TONGUE: *Pentstèmon hirsùtus.*
Figwort Family. May—July.

A plant 1–3 feet high, found in dry, rocky ground. Maine to Georgia, westward to Wisconsin and Missouri.

FLOWERS—Dull magenta or white, tubular, upper lip 2-lobed, lower lip 3–lobed, throat nearly closed by a hairy palate on the lower lip; flowers borne in loose spike.

LEAVES— Oblong to lance-shaped, slightly toothed, upper leaves stemless, lower leaves stemmed.

Digitized by Google

PAINTED CUP: Scarlet Painted Cup: Indian Paint-brush. *Castillèja coccínea.*

Figwort Family. May—July.

Found in wet meadows, moist, sandy soil. Massachusetts to Manitoba, south to Virginia, Tennessee and Texas.

FLOWERS—Small, greenish yellow, quite eclipsed by the 3-cleft floral leaves which are generally tipped with brilliant scarlet; flowers borne in clusters terminating stem 8–10 inches high.

LEAVES— Lower leaves oblong, clustered, undivided; upper ones generally deeply 3–5 cleft, stemless, stained on the tip with scarlet.

Digitized by Google

QUEEN OF THE PRAIRIE:

Filipéndula rùbra.

Rose Family. June—July.

A tall, smooth-stemmed, branching plant, 2–8 feet high, found in meadows, prairies. Pennsylvania to Michigan, Iowa, Kentucky and Georgia.

FLOWERS—Deep pink, fragrant, 5 divisions, borne in large clusters.

LEAVES— Compound 7–9-parted, terminal leaflet largest, lobed and toothed.

Digitized by Google

COMMON PASTURE, or DWARF, WILD ROSE: *Rósa virginiàna.*

Rose Family. June—July.

A shrub 1–5 feet high, found on margins of swamps, rocky shores. Newfoundland to New York and Eastern Pennsylvania.

FLOWERS—Crimson-pink, 5 divisions, outer divisions often with 1–2 small lobes.

LEAVES— Compound, usually 7 toothed leaflets, leaf-surface smooth and often shining above. Stipules (leafy formations at the bases of the leaves) more or less dilated. Prickles hooked.

SWAMP ROSE: *R. carolìna.*

A straight-stemmed shrub 2–7 feet high, found along borders of swamps and streams. Nova Scotia to Florida, westward to Minnesota and Mississippi. Leaflets 5–9, finely toothed, acute at each end, usually downy beneath. Stipules long and very narrow.

SMOOTH, EARLY or MEADOW ROSE: *R. blánda.*

A bush 2–4 feet high, found in rocky, moist ground. Newfoundland to New England and westward. Outer divisions of the flowers covered with bristly hairs. Leaflets 5–7, toothed. Stipules dilated, usually toothless. Prickles usually conspicuous by their absence.

Digitized by Google

PASTURE ROSE

R. hùmilis—Outer divisions of the flower always more or less lobed. Leaflets usually 7, smooth and often shining, toothed. Stipules usually narrow. Stems generally low, bearing straight, slender prickles. Nova Scotia to Florida, westward to Minnesota, Missouri and Louisiana.

SWEETBRIER; EGLANTINE:

R. rubiginòsa.

Readily known by the aromatic odor of its leaves when crushed. Introduced from Europe.

WILD BERGAMONT: *Monárda fistulòsa.*

Mint Family. July—Aug.

A plant 2–3 feet high, found in open woods, dry, rocky soil. Maine to Ontario, southward to Tennessee.

FLOWERS—Purplish, magenta, pink or whitish, tubular, 1–1½ inch long; upper lip erect, toothed, lower lip spreading, 3-lobed; clustered in a solitary, terminal head.

LEAVES— Opposite, lance-shaped, toothed, upper leaves whitish or the color of the flowers.

Digitized by Google

PURPLE CONE-FLOWER: *Braunèria purpùrea.*
Composite Family. July—Sept.

Found in rich soil. Western Pennsylvania and Virginia to Michigan, Iowa and southward.

FLOWER-HEADS—15–20 magenta colored, drooping rays, disk madder purple, terminating stem 2–3 feet high.

LEAVES— More or less toothed, lower ones 5-ribbed, long-stemmed, upper ones 3-veined, stemless.

B. pállida. Flowers similar to those of the preceding. Rays slender, drooping. Leaves lance-shaped, rough, narrowed at each end, 3-ribbed, toothless. Michigan and Illinois to Alabama and Texas.

CROSS-LEAVED MILKWORT: Marsh Milkwort. *Polýgala cruciàta.*
Milkwort Family. July—Sept.

A plant 3–12 inches high, found on margins of swamps, in low ground. Maine to South Carolina, and from Michigan to Minnesota and Nebraska.

FLOWERS—Magenta, white or greenish, tiny, borne in terminal, clover-like heads.

LEAVES— Generally in whorls of 4. Plant stems square, branching.

Digitized by Google

CORN COCKLE; CORN ROSE: Corn or Red Campion: Crown of the Field.

Agrostémma Githàgo.

Pink Family. July—Sept.

Found in grain fields, dry, waste places throughout the United States. Introduced from Europe.

FLOWERS—Magenta, 5 divisions, the narrow lobes of the outer divisions exceeding the inner ones in length; solitary, terminating stems 1–3 feet high.

LEAVES— Opposite, seated on stem, long, narrow, erect. Plant stem covered with fine, silky hairs.

SWAMP ROSE MALLOW; MALLOW ROSE.

Hibíscus Moscheùtos.

Mallow Family. July—Sept.

A shrub-like plant 4–6 feet high, found in marshes, on lake-shores, river-banks. Massachusetts and southward; usually near the coast; locally in the interior.

FLOWERS—Pink or white, 4–7 inches across, 5 divisions, conspicuously veined; flowers borne singly, or in scant clusters at the summit of the stems.

LEAVES— Egg-shaped, pointed, toothed, densely white-woolly beneath. Lower leaves and sometimes the upper ones 3-lobed.

Digitized by Google

PURPLE GERARDIA: Large Purple Gerardia.
Gerárdia purpùrea.

Figwort Family. Aug.—Sept.

Found in low fields, moist, sandy soil. Massachusetts to Florida and Texas near the coast, westward along the Great Lakes to Wisconsin.

FLOWERS—Purplish, magenta to whitish, cup-shaped, with 5 flaring lobes, downy.

LEAVES— Opposite, small, narrow. Plant stem somewhat angled, with spreading branches.

NOTES.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

IV

BLUE AND PURPLE

Digitized by Google

LIVERLEAF; HEPATICA: LIVERWORT: Round-lobed or Kidney Liverleaf: Noble Liverwort: Squirrel Cup. *Hepática tríloba.*

Crowfoot Family. Mch.—May.

Found in open woods. Nova Scotia to Florida, Missouri and Minnesota.

FLOWERS—Variable in color, blue-violet to white, 6–12 divisions, borne on hairy stems 4–6 inches high.

LEAVES— From the root, with 3 rounded lobes, green or reddish purple, new leaves appearing after the flowers bloom.

SHARP-LOBED LIVERLEAF: *H. acutíloba.* Differs from the preceding in that the 3 leaflets just beneath the flower, and lobes of the leaves are more acute at the tips. Leaves sometimes 5-lobed. More common westward.

Digitized by Google



Liverleaf.

Digitized by Google

BIRD-FOOT VIOLET: *Viola pedàta.*
Violet Family. Apr.—June.

Found in open, sunny fields and slopes. Massachusetts to Minnesota and southward.

FLOWERS—Largest of all the violets, blue-violet or lilac, orange center conspicuous, borne on stalk 4–10 inches high.

LEAVES— Variously divided and toothed.

NOTES.

Digitized by Google

Bird-foot Violet.

Digitized by Google

1. ROBIN'S PLANTAIN: Poor Robin's or Robert's Plantain: Blue Spring Daisy.
Erígeron pulchéllus.

Composite Family. Apr.—June.

An erect, hairy-stemmed plant 10–20 inches high, found in fields, copses, along roadsides. Maine to Ontario, Minnesota and southward.

FLOWER-HEADS—Rays lilac or pale violet, disk greenish yellow, borne in small terminal groups.

LEAVES— Lower leaves broad near the tip, slightly toothed, upper leaves lance-shaped.

2. WILD LUPINE: Old Maid's Bonnets: Wild Pea: Sun Dial. *Lupìnus perénnis.*

Pulse Family. May—June.

An erect, branching, leafy plant, found in dry, sandy places. New England to Minnesota, southward to the Gulf of Mexico.

FLOWERS—Violet-blue, blossoms pea-like in long clusters.

LEAVES— Compound, 7–11 long, narrow leaflets radiating from a common center.

Digitized by Google

1. Robin's Plantain. 2. Wild Lupine.

Digitized by Google

1. BLUETS; INNOCENCE: Quaker Ladies: Quaker Bonnets: Venus Pride: Houstonia.

Houstònia caerùlea.

Madder Family. Apr.—July.

A delicate little plant found in patches, almost carpeting moist, grassy places. Nova Scotia to Ontario and Wisconsin, southward to Georgia and Alabama.

FLOWERS—Pale violet, lilac or white, center yellow, about ½ inch across, 4-lobed, on slender stems 3–6 inches high.

LEAVES— Opposite, seated on stem, lower ones broader near the tip.

H. purpùrea is a southern species. Flowers deep or pale lilac, in small clusters. Leaves broad or lance-shaped, 3–5 ribbed. Plant stems smooth or hairy. Maryland to Iowa and southwest.

2. GILL-OVER-THE-GROUND: Ground Ivy or Joy: Creeping Charlie: Field Balm.

Népeta hederàcea.

Mint Family. Apr.—July.

A plant with creeping stem, 6–18 inches long, found in damp, shady places. Eastern half of United States and Canada, south to Kansas and Georgia. Naturalized from Europe.

FLOWERS—Small, pale purple, spotted darker near the throat; 2-lipped, upper lip 2-lobed, lower lip 3-lobed; growing from leaf angles.

LEAVES— Rounded, heart-shaped, scallop toothed.

Digitized by Google

1. Bluets. **2. Gill-over-the-Ground.**

Digitized by Google

LARGER BLUE FLAG; FLEUR-DE-LIS: Blue Iris: Flower-de-luce. *Iris versicolor.*
Iris Family. May—July.

Found in swamps, on margins of ponds and streams. Newfoundland to Manitoba and southward.

FLOWERS—Violet-blue, the 3 outer divisions recurved, yellow at the base, veined with purple; 1 or more flowers borne on stem 2–3 feet high.

LEAVES— Erect, sword-shaped.

SLENDER BLUE FLAG: *I. prismática.*

Flowers smaller than those of the preceding, borne on slender stems. Leaves narrow, grass-like. Marshes near the coast. Nova Scotia to Georgia.

NOTES.

Digitized by Google

Larger Blue Flag.

Digitized by Google

COW VETCH; TARE: Blue or Tufted Vetch: Cat Peas: Tinegrass. *Vícia Crácca.*

Pulse Family. June—Aug.

Found in fields, waste land. Newfoundland to New Jersey, westward to Kentucky, Iowa and Minnesota.

FLOWERS—Small, violet-blue, closely set on a 1-sided spike. Fruit a small pod.

LEAVES— Tendril-bearing, divided into 18–24 narrow leaflets with bristle-like points.

AMERICAN VETCH: Tare or Pea Vine. *V. americàna.*

Flowers 3–9, slightly larger than those of the preceding. Leaflets 8–14, blunt; distinctly veined. Found in moist places. New York to Virginia and westward.

NOTES.

Digitized by Google



Cow Vetch.

Digitized by Google

BITTERSWEET; NIGHTSHADE: Blue Bindweed: Felonwort: Scarlet or Snakeberry: Poison Flower: Woody Nightshade.

Solànum Dulcamàra.

Nightshade Family. June—Sept.

A climbing or straggling, shrubby plant, stems 2–8 feet long, found in moist thickets. Eastern United States. Naturalized from Europe.

FLOWERS—Violet or purple, with yellow conic center, 5 reflexed divisions, in loose clusters. Berries inedible, successively green, yellow, orange, then red; appear at the same time with flowers.

LEAVES— Toothless, variable, some bearing 2 leaflets at the base.

NOTES.

Digitized by Google

Bittersweet.

Digitized by Google

1. SELF-HEAL; HEAL-ALL: Blue Curls: Heart-of-the-Earth: Brunella.

Prunélla vulgàris.

Mint Family. June—Sept.

Found in fields, waste places, along roadsides, throughout the United States. Naturalized from Europe.

FLOWERS—Tiny, purple, varying to white, lower lip slightly fringed, borne in a dense head, terminating stem 6–12 inches high.

LEAVES— Opposite, oblong, narrowed toward the tip, slightly toothed.

2. BLUE-EYED GRASS: Eye-bright: Blue Star. *Sisyrínchium angustifòlium.*

Iris Family. May—July.

Found in moist fields, meadows. Newfoundland to the far West, southward to Virginia.

FLOWERS—Deep violet-blue with a yellow center, the 6 divisions tipped with a point; flowers borne erect on flat stems 6–14 inches high.

LEAVES— Grass-like.

Digitized by Google

1. Self-heal. 2. Blue-eyed Grass.

Digitized by Google

1. BELLFLOWER: Rampion, European, or Creeping Bellflower. *Campánula rapunculoìdes.*
Bluebell Family. July—Aug.
A sturdy plant 1–3 feet high, found along roadsides and in waste places. New Brunswick to Ontario, Pennsylvania and Ohio. Introduced from Europe and Asia. Escaped from gardens.
FLOWERS—Purple-blue, bell-shaped, 5-lobed, hanging down mostly on one side of the stem.
LEAVES— Lower leaves arrowhead-shaped, with a heart-shaped base; upper leaves broad lance-shaped, short-stemmed, toothed.

2. HAREBELL; BLUEBELL: Bluebells of Scotland: Lady's Thimble.
Campánula rotundifòlia.
Bluebell Family. June—Sept.
Found on dry or moist rocky cliffs, in meadows, along margins of ponds. Arctic regions south to New Jersey and westward.
FLOWERS—Violet-blue, bell-shaped, 5-lobed, terminating exquisitely slender stalks 6–18 inches high.
LEAVES— Lower leaves nearly round, usually withered before flowering time; stem leaves narrow, seated on stem.

Digitized by Google

1. Bellflower. **2. Harebell.**

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

1. PALE SPIKED LOBELIA: *Lobèlia spicàta.*

Lobelia Family. July—Aug.

Found in dry, sandy soil, meadows. Prince Edward Island to Ontario, westward and southwest.

FLOWERS—Pale bluish violet, upper lip of 2 narrow, erect lobes, lower lip spreading, 3-lobed; flowers borne in loose spikes, terminating stem 1–4 feet high.

LEAVES— Lower leaves oblong, obtuse, upper leaves narrow, slightly toothed.

2. MOTHERWORT: *Leonùrus Cardìaca.*

Mint Family. June—Aug.

An erect, branched plant 2–4 feet high, found in waste places, generally near dwellings. Nova Scotia to North Carolina, west to Minnesota and Nebraska. Naturalized from Europe.

FLOWERS—Pale lilac or white, small, upper lip bearded, clustered in the axils of the upper leaves.

LEAVES— Opposite, long-stemmed, lower leaves variously cut and toothed, upper leaves 3-cleft.

Digitized by Google

2

1. Pale Spiked Lobelia. **2. Motherwort.**

Digitized by Google

1. PICKEREL-WEED: *Pontedèria cordàta.*
Pickerel-weed Family. July—Sept.

An erect, stout, fleshy plant 1–3 feet high, found in shallow water. Nova Scotia to Ontario, Minnesota and Texas.

FLOWERS—Violet-blue, 2-lipped, upper lip 3-lobed, with 2 yellowish spots within; lower lip of 3 spreading lobes; flowers borne on thick spike.

LEAVES— One large leaf shaped like an arrowhead, is borne midway of the flower stalk.

2. MONKEY FLOWER: *Mímulus ríngens.*
Figwort Family. June—Sept.

A square-stemmed plant 1–3 feet high, found beside brooks and streams, swamps. New Brunswick to Manitoba and southward.

FLOWERS—Purple, with 2 yellow spots near the throat; 2-lipped, upper lip 2-lobed, erect; lower lip 3-lobed, spreading; flowers borne on slender stalks from the leaf angles.

LEAVES— Opposite, oblong or lance-shaped, toothed, clasping stem by a heart-shaped base.

Digitized by Google

1. Pickerel-weed. **2. Monkey Flower.**

Digitized by Google

1. MAD-DOG SKULLCAP: Helmet Flower: Madweed: Hoodwort. *Scutellària lateriflòra.*
Mint Family. July—Sept.
A plant 1–2 feet high, found in damp, shady places throughout the United States and northward.
FLOWERS—Blue or purple, varying to whitish, small, about ¼ inch long, growing in 1-sided spikes from the angles of the upper leaves.
LEAVES— Opposite, pointed, rounded at the base, coarsely toothed. 1–2 feet high.

2. SKULLCAP: Larger or Hyssop Skullcap. *Scutellària integrifòlia.*
Mint Family. May—Aug.
Found in borders of fields and woods. Massachusetts to Florida and Texas, chiefly near the coast.
FLOWERS—Blue, about 1 inch long, opposite each other at the top of a stem 6–20 inches high.
LEAVES— Oblong or narrower, mostly toothless. Plant stem and leaves covered with a fine down.

Digitized by Google

1. Mad-dog Skullcap. 2. Skullcap.

Digitized by Google

BLUE VERVAIN: Wild Hyssop: Simpler's Joy.
Verbèna hastàta.

Vervain Family. July—Sept.

A plant 3–7 feet high, found in fields, waste places, along roadsides. Common in the United States and Canada.

FLOWERS—Purple or violet, small, 5-lobed, blossoming in circles from the base of the flower spikes upward.

LEAVES— Opposite, stemmed, lance-shaped, toothed, rough; lower leaves more or less lobed.

NOTES.

Digitized by Google



Blue Vervain.

Digitized by Google

1. WILD MINT:

Méntha arvénsis var. canadénsis.

Mint Family. July—Sept.

Found in wet places. New Brunswick to California.

FLOWERS—Purplish to white, tiny in whorls at the angles of the leaves.

LEAVES— Tapering from the center toward each end, toothed, hairy throughout, aromatic odor like pennyroyal. Plant stems branching, 10–24 inches high.

2. SPEARMINT: Garden or Mackerel Mint.

Méntha spicàta.

Mint Family. July—Sept.

Found in wet places. Eastern half of United States and Canada. Naturalized from Europe.

FLOWERS—Purplish, variable in depth of color, in narrow, acute, usually interrupted terminal spikes.

LEAVES— Aromatic, opposite, narrowly oblong, acute, sharply toothed, stemless or short-stemmed.

3. PEPPERMINT: *M. piperita.*

Flowers purplish, in dense or interrupted, blunt, terminal spikes. Leaves egg-shaped, smooth, toothed, acute, very pungent tasting. Plant stem purplish, 18–36 inches high. Naturalized from Europe.

Digitized by Google



1. Wild Mint. 3. Peppermint.

2. Spearmint.

Digitized by Google

IRONWEED: Flat Top.

Vernònia noveboracénsis.

Composite Family. July—Sept.

Found in moist soil, meadows. Massachusetts to Virginia and Mississippi; usually near the coast.

FLOWER-HEADS—Reddish purple, the brownish purple scales bristle tipped, in broad, flat clusters terminating stem 3–9 feet high.

LEAVES— Long lance-shaped, toothed.

NOTES.

Digitized by Google

Ironweed.

Digitized by Google

1. CHICORY; SUCCORY: Blue Sailors: Bunk: *Cichòrium Intybus.*

Composite Family. July—Oct.

Found in fields, waste places, along roadsides. Newfoundland to Minnesota and southward. Naturalized from Europe.

FLOWER-HEADS—Blue, violet-blue, lavender, rays finely toothed on the tips; flowers borne at intervals on branching stems 1–3 feet high.

LEAVES— Oblong or lance-shaped, coarsely toothed, partly clasping.

2. BLAZING STAR: Button Snakeroot: Gay Feather. *Liàtris scariòsa.*

Composite Family. Aug.—Sept.

Found in fields, along roadsides, dry soil. Maine to Ontario, Nebraska and southward.

FLOWER-HEADS—Variable purple, borne in a spike terminating stem 2–6 feet high.

LEAVES— Narrow lance-shaped.

Digitized by Google

1. Chicory. 2. Blazing Star.

Digitized by Google

NEW ENGLAND ASTER: Starwort.

Aster nòvae-ángliae.

Composite Family. Aug.—Oct.

Most striking of the asters; found in swamps, roadsides, moist ground. Maine, westward and southwest. Like the golden-rods, the number of different species of asters, with their intergrading forms, is so great as to render their identification difficult, even for the experienced botanist.

FLOWER-HEADS—Rays 40–50, light purple or violet, disk yellow.

LEAVES— Lance-shaped, toothless, clasping stem by a broad base. Plant stem stout, branched, rough.

Digitized by Google

New England Aster.

Digitized by Google

1. FRINGED GENTIAN: *Gentiàna crinìta.*
Gentian Family. Sept.—Oct.

Found in moist meadows, woods. Maine to Quebec, south to Georgia and westward.

FLOWERS—Pale or deep violet-blue, erect, tubular, with 4 spreading, fringed lobes.

LEAVES— Opposite, acute at the tip, widening to a rounded base, seated on stem. Plant stem branching, erect, each bearing a single terminal flower, 1–3 feet high.

2. CLOSED GENTIAN: Blind or Bottle Gentian. *Gentiàna Andréwsii.*
Gentian Family. Aug.—Oct.

A plant 1–2 feet high, found in woodland borders, moist soil. Maine to Manitoba and southward.

FLOWERS—Tightly closed, deep violet blue at the apex, white at the base, in terminal clusters from the leaf angles.

LEAVES— Opposite, smooth, pointed at the tip, narrowed at the base.

Digitized by Google

1. Fringed Gentian. **2. Closed Gentian.**

Digitized by Google

Digitized by Google

NOTES.

Digitized by Google

VIRGINIAN COWSLIP: Bluebells: Tree or Smooth Lungwort. *Mertènsia virgínica.*
Borage Family. Apr.—May.

A plant 1–2 feet high, found in low meadows, along streams. New York and Ontario to Nebraska, and southward.

FLOWERS—Pinkish in the bud, afterward violet, trumpet-shaped, 5-lobed, about 1 inch long, borne in loose clusters which are rolled up at the tip and straighten as the flowers expand.

LEAVES— Veiny, toothless, upper leaves seated on stem, lower ones long-stemmed.

EARLY BLUE VIOLET: *Vìola palmàta.*
Violet Family. Apr.—May.

Found in dry ground, chiefly woodlands. Massachusetts to Minnesota, and southward.

FLOWERS—Light or deep violet.

LEAVES— Heart-shaped, first leaves of spring sometimes undivided, later ones 5–9-lobed, the segments variously toothed or cleft, middle segment usually the largest.

Digitized by Google

ARROW-LEAVED VIOLET:

Viola sagittàta.

Violet Family. Apr.—May.

Plant 2–8 inches high; found in wet weadows, dry borders. Massachusetts to Minnesota and southward.

FLOWERS—Light or deep violet, upper and side divisions bearded, lower one veined.

LEAVES— Arrowhead-shaped, often toothless above the middle, slightly lobed below.

DOG VIOLET: Running Violet. *Viola conspérsa.*

Violet Family. Apr.—June.

Plant 2–6 inches high, found in low, shaded ground. Eastern Quebec to Minnesota and southward.

FLOWERS—Pale purple, sometimes white, raised above the leaves on stems from the leaf angles, pansy-fashion.

LEAVES— Round heart-shaped, with barely noticeable scallops. Plant stems ascending.

Digitized by Google

COMMON VIOLET: Purple, Meadow or Hooded Blue Violet. *Viola papilionàcea.*
Violet Family. Apr.—June.
Plant 3–7 inches high, found in low ground. Massachusetts to Minnesota and southward.
FLOWERS—Light purple, variable, divisions yellowish or white at the base.
LEAVES— Heart-shaped, scallop-toothed.

WILD BLUE PHLOX: *Phlóx divaricàta.*
Polemonium Family. May—June.
Plant 1–2 feet high, easily recognized by its likeness to the cultivated Phloxes; found in moist, thin woods. Western Quebec to Minnesota and southward.
FLOWERS—Pale violet or lilac, in loose, spreading clusters, 5 lobes, notched at the tips.
LEAVES— Those of flowering stem opposite, oblong, tapering to a point; those of sterile shoots oblong or egg-shaped, not pointed. Plant stem finely coated with rather sticky hairs.

Digitized by Google

FORGET-ME-NOT: Mouse-ear: Snake Grass: Scorpion Grass: Love Me: *Myosòtis scorpioìdes.*
Borage Family. May—July.
Escaped from gardens; a plant with leafy, branching, half-reclining stems 6–15 inches high, found in wet ground. Newfoundland to Pennsylvania and beyond. Naturalized from Europe.
FLOWERS—Small, light blue, with a golden eye, 5-lobed, in small clusters.
LEAVES— Oblong or lance-shaped, seated on stem; leaves and stem hairy.

SMALLER FORGET-ME-NOT: *M. láxa.*
A native species, found in wet ground from Newfoundland to Ontario and southward.
FLOWERS—Light blue, on long stems, loosely clustered; lobes of the flower cup as long as the flower tube.
LEAVES— Blunt, oblong.

SPIDERWORT: Widow's or Job's Tears. *Tradescántia virginiàna.*
Spiderwort Family. May—Aug.
Found in rich, moist woods, thickets. Connecticut to South Carolina; also introduced northwestward.
FLOWERS—Purplish blue, 1–2 inches broad, clustered and seated upon the end of stem between blade-like leaflets, 3 parted.
LEAVES— Long, narrow, sheathing at the base. Plant stem erect, leafy, mucilaginous, 8–36 inches high.

Digitized by Google

BEACH PEA: Seaside, Sea, or Everlasting Pea. *Láthyrus marítimus.*

Pulse Family. May—Aug.

Found chiefly near seashores from New Jersey and Oregon to the Arctic Sea; also shores of Great Lakes.

FLOWERS—Purple, about 1 inch long or less, 5–12 in short clusters on stalk springing from the leaf angles.

LEAVES— Compound, consisting of 6–10 leaflets. A pair of arrowhead-shaped leaflets are borne at the base of the compound leaves. Plant stem stout, reclining, 1–2 feet long.

COMMON SPEEDWELL: Fluellin: Paul's Betony: Ground-hele. *Verónica officinàlis.*

Figwort Family. May—Aug.

A low growing plant, with downy, prostrate stem 3–10 inches long, found in dry fields, open woods. Newfoundland to Ontario, Michigan, and southward.

FLOWERS—Small, light lavender in color, with darker lines, 4 unequal divisions, borne in slender spikes from the leaf angles, held erect from the hairy, prostrate stem.

LEAVES— Oval, toothed, rounded at the tip, tapering at the base into a short stem.

Digitized by Google

AMERICAN BROOKLIME:

Verónica americàna.

Figwort Family. May—Sept.

A smooth-stemmed plant, 6–15 inches high, its stem lying partly on the ground, rooting from lower joints; found on borders of brooks, ponds, damp places. Newfoundland to Alaska and southward to Virginia.

FLOWERS—Small, quickly fading; in color lavender-blue, with slightly darker markings, 4-lobed, the lower lobe narrower than the others; borne in loose, terminal clusters.

LEAVES— Opposite, long oval or oblong lance-shaped, toothed, short-stemmed.

VIRGINIA, or COMMON, DAY-FLOWER:

Commelìna virgínica.

Spiderwort Family. June—Sept.

A plant with smooth, branched stems 18–36 inches high, found in moist, shady places. New York to Florida, west to Michigan, Kansas and Texas.

FLOWERS—Violet-blue, about 1 inch broad, of 3 divisions, one of which is small and inconspicuous.

LEAVES— Lance-shaped, sheathing. A small leaf more or less encloses flower cluster.

C. hirtélla is similar to the preceding, but may be distinguished by its leaves, which are brown-bearded where they sheath the flower stem.

Digitized by Google

BLUE TOADFLAX: Wild Toadflax: Blue Linaria. *Linària canadénsis.*

Figwort Family. June—Sept.

A slender plant, 5–30 inches high, found in dry, sandy places. New England westward and southwest.

FLOWERS—Pale violet or lavender, small, in slender spikes; upper lip with 2 acute divisions, lower lip larger, 3-lobed, white at the throat.

LEAVES— Narrow, toothless.

VIPER'S BUGLOSS; BLUE-WEED: Viper's Herb or Grass: Blue Thistle: Snake-flower: Blue Devil. *Èchium vulgàre.*

Borage Family. June—Sept.

A plant 12–30 inches high, with bristly, hairy, spotted stem, found in fields, waste places, along roadsides. Naturalized from Europe.

FLOWERS—Variable purple, pink in the bud, 5 unequal lobes, borne on a 1-sided spike, which is coiled at first, unrolling as the flowers expand.

LEAVES— Lance-shaped, hairy, mostly stemless.

Digitized by Google

TALL WILD LARKSPUR:

Delphínium exaltàtum.

Crowfoot Family. July—Aug.

A plant 2–6 feet high, found in rich woods. Pennsylvania to Minnesota, Nebraska and southward.

FLOWERS—Light purple, irregular, borne in long, slender spikes.

LEAVES— Deeply 3–5-cleft, divisions narrowly wedge-shaped, 3-cleft at the apex.

DWARF LARKSPUR: Stagger Weed.

D. tricórne.

A plant 2–3 feet high, appearing earlier in the season; leaves deeply 5-parted, their divisions unequally 3–5-cleft; root a tuberous cluster.

PURPLE ORCHIS: Fringeless Purple Orchis.

Habenària peramoèna.

Orchis Family. July—Aug.

Found in moist meadows. Pennsylvania and New Jersey to Illinois, and southward.

FLOWERS—Purple, clustered in dense spike terminating stem 12–30 inches high; lower lip fan-shaped, toothed.

LEAVES— Oblong to lance-shaped.

Digitized by Google

SEA LAVENDER; MARSH ROSEMARY: Canker-root: Ink-root. *Limònium carolinià̀num.* Leadwort Family. July—Sept.

A plant 1–2 feet high, with much-branched stems, found in salt marshes. Labrador to Texas.

FLOWERS—Lavender, very tiny, borne in sprays on upper side of branches.

LEAVES— From the root, oblong, or blunt lance-shaped, midrib prominent.

NOTES.

Digitized by Google

NOTES.

Digitized by Google

Digitized by Google

V

MISCELLANEOUS

Digitized by Google

SKUNK CABBAGE: Swamp Cabbage.

Symplocárpus foétidus.

Arum Family. Mch.—Apr.

An unmistakably ill-scented plant, found in low, wet ground. Nova Scotia to North Carolina, westward to Ontario, Minnesota and Iowa.

FLOWERS—Minute, scattered over a thick, fleshy club within its flower envelope, which is variously streaked with greenish yellow and madder purple, and appears just above the ground.

LEAVES— From the base, large, often 1 foot across, strongly veined.

NOTES.

Digitized by Google



Skunk Cabbage.

Digitized by Google

WILD GINGER: Canada Snakeroot: Asarabacca.
Asarum canadénse.

Birthwort Family. Apr.—May.

Found in rich woods. North Carolina, Missouri and Kansas, northward to New Brunswick and Manitoba.

FLOWERS—Brownish purple, with 3 long-pointed divisions, borne on short stem close to the ground, often hidden under the leaves.

LEAVES— 2, broadly heart-shaped, leaf stems hairy, 6–12 inches high, from aromatic rootstock.

NOTES.

Digitized by Google



Wild Ginger.

Digitized by Google

JACK-IN-THE-PULPIT; INDIAN TURNIP:
Arisaèma triphýllum.

Arum Family. Apr.—June.

A plant 10–30 inches high, springing from an acrid bulb, found in moist, shady woods. Nova Scotia, westward to Minnesota, and southward to the Gulf States.

FLOWERS—Minute, clustered at the base of the smooth club within the flower envelope; latter pale green or marked with purplish stripes, variable in depth of color. Berries red, clustered in a head.

LEAVES— 1–2, long-stemmed, smooth, divided into 3 large leaflets.

NOTES.

Digitized by Google



Jack-in-the-Pulpit.

Digitized by Google

1. BRANCHING BUR-REED: *Spargànium americànum.*

Bur-reed Family. June—Aug.

Plant 1–2 feet high, stem having 1–2 branches, found on borders of ponds and streams. Newfoundland to Minnesota, southward to Missouri and Florida.

FLOWERS—Of 2 kinds, the upper ones in downy heads, lower ones round, bur-like, borne on stalks from the leaf angles.

LEAVES— Long, narrow.

2. SWEET FLAG: *Acorus Cálamus.*

Arum Family. June—Aug.

Found in wet, swampy places. Maine, south and west to Minnesota, Iowa, and Kansas.

FLOWERS—Tiny, compactly covering a long, slender, greenish club, which grows from the side of a two-edged stem.

LEAVES— Long, narrow, sword-like. Rootstock aromatic.

Digitized by Google



1. Branching Bur-reed.

2. Sweet Flag.

Digitized by Google

1. PINESAP; FALSE BEECHDROPS: Yellow Bird's-nest. *Monótropa Hypópitys.*

Heath Family. June—Sept.

Found in dry woods. Florida and Arizona, northward into British Possessions.

FLOWERS—Crimson and yellow, somewhat variable, about ½ inch long, slightly fragrant, borne in drooping clusters (which become erect later), terminating stem 4–12 inches high.

LEAVES— None. Stems clustered, scaly, arising from a mass of fibrous roots.

2. BEECHDROPS: Cancer-root. *Epifàgus virginiàna.*

Broom-rape Family. Aug.—Oct.

Found usually under beech trees. New Brunswick to Ontario, Wisconsin, and southward.

FLOWERS—Light yellow or brownish, purple-striped, scattered along the branches.

LEAVES— None. Plant stem brownish or reddish tinged, arising from brittle, fibrous roots, 6–20 inches high.

Digitized by Google

1. Pinesap. 2. Beechdrops.

Digitized by Google

1. LARGE CORAL ROOT:
Corallorrhiza maculàta.

Orchis Family. July—Aug.

Found in dry woods. Nova Scotia, westward to British Columbia, south to Florida, Missouri, and California.

FLOWERS—Light yellow, more or less purplish-stained; lower lip white with purplish spots and lines, 3-lobed; borne on spike terminating stem 8–20 inches high.

LEAVES— None. Plant stem madder-purple or yellowish.

2. RATTLESNAKE-ROOT: Wild Lettuce: Canker-weed: Lion's Foot. *Prenánthes altíssima.*

Composite Family. July—Sept.

Plant 3–6 feet high, found in woodlands, thickets. Newfoundland to Manitoba, southward to Georgia and Tennessee.

FLOWER-HEADS—Inner divisions inconspicuous, whitish, pendulous in loose terminal cluster, with small clusters from the leaf angles.

LEAVES— Long-stemmed, variously shaped, toothed, and cleft.

Digitized by Google



1. Coral Root. 2. Rattlesnake-root.

Digitized by Google

GROUNDNUT: Wild Bean. *Apios tuberòsa.*
Pulse Family. July—Sept.

A climbing vine, 3–6 feet high, found in low, moist ground. New Brunswick to Minnesota, south to Florida, Louisiana, and Kansas.

FLOWERS—Maroon and pale brown-lilac, fragrant, borne in short, thick clusters from the leaf angles.

LEAVES— Compound, 3–7 oval leaflets.

Rootstock bears edible, tuberous enlargements.

NOTES.

Digitized by Google

Groundnut.

Digitized by Google

Digitized by Google

NOTES.

Digitized by Google

EARLY MEADOW RUE:

Thalictrum dioicum.

Crowfoot Family. Apr.—May.

Plant 1–2 feet high, found in thin woods. Maine, westward and southward.

FLOWERS—Small, tassel-like, in airy clusters, green, or purplish in effect.

LEAVES— Compound, leaflets small, lustreless, 3–7-lobed.

BLUE COHOSH: Pappoose Root.

Caulophýllum thalictroìdes.

Barberry Family. Apr.—May.

Plant 1–3 feet high, found in rich woods. New Brunswick to Manitoba and southward.

FLOWERS—Greenish yellow, more or less tinged with the bluish color of the plant; flowers about ½ inch across, 6 divisions, borne in small clusters, which appear before the leaves are fully expanded. Seeds blue.

LEAVES— Compound, leaflets lobed. Whole plant covered with a whitish bloom when young.

Digitized by Google

COMMON ALUM ROOT:

Heuchèra americàna.

Saxifrage Family. May—Aug.

Found in dry and rocky woods. Connecticut to North Carolina, westward to Minnesota, Kansas, and Mississippi.

FLOWERS—Greenish or purplish, 5-cleft, with several orange-tipped filaments protruding; borne in a long, narrow cluster on a more or less hairy stem, 2–3 feet high.

LEAVES— Heart-shaped, with rounded lobes and teeth.

NOTES.

Digitized by Google

LARGE TWAYBLADE: *Liparis liliifòlia.*
Orchis Family. June—July.

Found in woodlands. New Hampshire to Minnesota, Missouri, and Alabama.

FLOWERS—Madder-purple or greenish, lip broad, translucent, other divisions exceedingly narrow; 5–15 flowers borne on stem 4–9 inches high.

LEAVES— 2, from the base, broad, shining.

NOTES.

Digitized by Google

CUT-TAIL FLAG: *Typha latifòlia.*

Bur-reed Family. June—July.

Found in swamps throughout temperate North America.

FLOWERS—Borne in a dense, cylindrical head terminating stem 4–8 feet high.

LEAVES— Long, narrow, usually exceeding the flower stem in height.

NARROW-LEAVED CAT-TAIL: *T. angustifòlia.*

Leaves narrower than those of the preceding, and the 2 parts of the flower-head are usually separated by a short gap. Plant 4–9 feet high. Maine to North Carolina and westward, mainly near the coast.

NOTES.

Digitized by Google

NOTES.

Digitized by Google

NOTES.

Digitized by Google

Digitized by Google

INDEX

	PAGE
Achillea Millefolium	58
Acorus Calamus	328
Actaea alba	14
Actaea rubra	14
Adder's Tongue, White	96
Adder's Tongue, Yellow	110
Agrimonia gryposepala	138
Agrimony	138
Agrostemma Githago	265
Alder, White	40
Alisma Plantago-aquatica	82
Allium tricoccum	46
Alum Root, Common	339
Anaphalis margaritacea	86
Anemone quinquefolia	18
Anemone, Rue	18
Anemone, Fall or Summer	44
Anemone virginiana	44
Anemone, Wood	18
Anemonella thalictroides	18
Antennaria plantaginifolia	12
Anthemis Cotula	54
Apios tuberosa	334
Apocynum androsaemifolium	226
Aquilegia canadensis	194
Aralia hispida	36
Aralia nudicaulis	20
Arbutus, Trailing	94
Arethusa bulbosa	190
Arisaema triphyllum	326
Arrow-head	72
Arum, Green Arrow	24
Arum, Water	26
Asarum canadense	324

Digitized by Google

PAGE

Asclepias incarnata 242
Asclepias phytolaccoides 102
Asclepias purpurascens 236
Asclepias quadrifolia 242
Asclepias syriaca 236
Asclepias tuberosa 183
Aster, Golden 185
Aster, New England 304
Aster novae-angliae 304
Azalea, Clammy 100
Azalea, Wild or Pink 208

Balmony 70
Balsam-apple, Wild 76
Baneberry, Red 14
Baneberry, White 14
Barbarea vulgaris 150
Beard-tongue, Hairy 259
Bee Balm 246
Bedstraw, Rough 72
Beechdrops 330
Beechdrops, False 330
Bellflower 286
Bellwort 122
Bellwort, Large-flowered 122
Bergamot, Wild 263
Betony, Wood 216
Bidens laevis 176
Bindweed, Field 238
Bindweed, Hedge 238
Bird's Nest 88
Birthroot 192
Bishop's Cap 10
Bitter Bloom 240
Bittersweet 282
Black-eyed Susan 142
Blazing Star 302
Bloodroot 2
Bluebell 286
Bluets 276
Blue-weed 316
Boneset 68
Bouncing Bet 216
Brassica nigra 150

Digitized by Google

PAGE

Brauneria pallida 264
Brauneria purpurea 264
Brooklime, American 315
Bugloss, Viper's 316
Bunchberry 54
Bur-marigold, Larger 176
Bur-reed, Branching 328
Butter-and-Eggs 160
Butterfly Weed 183
Buttonbush 90

Calico Bush 212
Calla, Marsh 26
Calla palustris 26
Calopogon pulchellus 222
Caltha palustris 112
Campanula rapunculoides 286
Campanula rotundifolia 286
Campion, Bladder 50
Campion, White 50
Cancer-root 330
Canker-root 30
Cardinal Flower 244
Carrion-flower 130
Carrot, Wild 88
Cassia Chamaecrista 166
Cassia marilandica 166
Castalia odorata 103
Castilleja coccinea 260
Caulophyllum thalictroides 338
Ceanothus americanus 100
Celandine 138
Cephalanthus occidentalis 90
Cerastium arvense 44
Chamomile 54
Checkerberry 104
Cheeses 52
Chelidonium majus 138
Chelone glabra 70
Chickweed, Field 44
Chicory 302
Chimaphila maculata 234
Chimaphila umbellata 234
Chiogenes hispidula 99

Digitized by Google

PAGE

Chrysopsis mariana 185
Cichorium Intybus 302
Cicuta maculata 102
Cimicifuga racemosa 100
Cinquefoil 114
Claytonia caroliniana 188
Claytonia virginica 188
Clematis virginiana 60
Clethra alnifolia 40
Clintonia borealis 118
Clintonia, Yellow 118
Clover, Sweet 86–181
Clover, Rabbit's-foot 252
Clover, Yellow or Hop 148
Cohosh, Black 100
Cohosh, Blue 338
Collinsonia canadensis 184
Columbine, Wild 194
Commelina virginica 315
Cone-flower 142
Cone-flower, Purple 142–264
Cone-flower, Tall 142
Convolvulus arvensis 238
Convolvulus sepium 238
Coptis trifolia 80
Coral Root, Large 332
Corallorrhiza maculata 332
Corn Cockle 265
Cornel, Dwarf 54
Corn Rose 265
Cornus canadensis 54
Corpse plant 46
Corydalis aurea 179
Corydalis, Golden 179
Corydalis, Pale 218
Corydalis sempervirens 218
Cowbane, Spotted 102
Cowslip, American 190
Cowslip, Virginian 310
Cranesbill, Wild 198
Cress, Water 99
Cress, Winter 150
Crinkle-root 6
Cucumber-root, Indian 120

Digitized by Google

PAGE

Culver's Root 105
Cuscuta Gronovii 84
Cypripedium acaule 204
Cypripedium hirsutum 101
Cypripedium parviflorum 126
Cypripedium parviflorum var. pubescens . . . 126

Dalibarda repens 103
Dandelion, Fall 166
Datura Stramonium 105
Daucus Carota 88
Day-flower, Virginia 315
Deer-grass 246
Delphinium exaltatum 317
Delphinium tricorne 317
Dentaria diphylla 6
Dentaria laciniata 6
Devil's Paint-brush 172
Dianthus Armeria 196
Dianthus deltoides 196
Dicentra canadensis 4
Dicentra Cucullaria 4
Dodder, Common 84
Dodecatheon Meadia 190
Dogbane, Spreading 226
Drosera longifolia 66
Drosera rotundifolia 66
Dutchman's Breeches 4

Echinocystis lobata 76
Echium vulgare 316
Eglantine 263
Elder, Black-berried 40
Elder, Common 40
Elder, Red-berried 40
Elecampane 164
Epifagus virginiana 330
Epigaea repens 94
Epilobium angustifolium 244
Epipactis pubescens 62
Erigeron annuus 56
Erigeron pulchellus 274
Erigeron ramosus 56
Erythronium albidum 96

Digitized by Google

PAGE

Erythronium americanum 110
Eupatorium perfoliatum 68
Eupatorium purpureum 250
Eupatorium urticaefolium 68
Everlasting, Early Spring 12
Everlasting, Pearly 86
Everlasting, Plantain-leaved 12

Filipendula rubra 261
Fireweed 244
Flag, Cat-tail 341
Flag, Larger Blue 278
Flag, Narrow-leaved Cat-tail 341
Flag, Slender Blue 278
Flag, Sweet 328
Fleabane, Daisy 56
Fleur-de-lis 278
Flower-de-luce 278
Foam-flower 10
Forget-me-not 313
Forget-me-not, Smaller 313
Foxglove, Downy False 162
Foxglove, Fern-leaved False 185
Foxglove, Smooth False 162
Frostweed 182

Galium asprellum 72
Gaultheria procumbens 104
Gentian, Closed 306
Gentian, Fringed 306
Gentiana Andrewsii 306
Gentiana crinita 306
Geranium maculatum 198
Geranium Robertianum 198
Geranium, Wild 198
Gerardia flava 162
Gerardia pedicularia 185
Gerardia purpurea 266
Gerardia, Purple 266
Gerardia virginica 162
Gill-over-the-Ground 276
Ginger, Wild 324
Ginseng 183
Ginseng, Dwarf 20

Digitized by Google

PAGE

Globe-flower 90
Golden Alexanders 116
Golden Club 180
Golden-rod 185
Goldthread 30
Grass, Blue-eyed 284
Grass of Parnassus 80
Groundnut 334

Habenaria blephariglottis 104
Habenaria bracteata 42
Habenaria ciliaris 162
Habenaria fimbriata 234
Habenaria Hookeri 64
Habenaria hyperborea 42
Habenaria lacera 62
Habenaria leucophaea 104
Habenaria orbiculata 64
Habenaria peramoena 317
Habenaria psycodes 234
Hardhack 228
Harebell 286
Hawkweed, Canada 172
Hawkweed, Hairy 174
Hawkweed, Orange or Tawny 172
Hawkweed, Rough 174
Heal-all 284
Helenium autumnale 184
Helianthemum canadense 182
Helianthus decapetalus 176
Hellebore, American White 181
Hemlock, Water 102
Hepatica acutiloba 270
Hepatica triloba 270
Herb Robert 198
Heuchera americana 339
Hibiscus Moscheutos 265
Hieracium aurantiacum 172
Hieracium canadense 172
Hieracium Gronovii 174
Hieracium scabrum 174
Hieracium venosum 174
Honeysuckle 194
Honeysuckle, White Swamp 100

Digitized by Google

PAGE

Horse Balm 184
Houstonia caerulea 276
Houstonia purpurea 276
Hypericum perforatum 158
Hypoxis hirsuta 114

Impatiens biflora 156
Impatiens pallida 156
Indian Hemp 226
Indian Pipe 46
Indian Poke 181
Indian Turnip 326
Innocence 276
Inula Helenium 164
Iris prismatica 278
Iris versicolor 278
Ironweed 300
Ivy, Poison 34

Jack-in-the-Pulpit 326
Jamestown or Jimson Weed 105
Jeffersonia diphylla 96
Jewel-weed 156
Joe-Pye Weed 250

Kalmia angustifolia 220
Kalmia latifolia 212
Kalmia polifolia 220
Knotweed, Pink 248

Lactuca canadensis 146
Ladies' Tresses 66
Ladies' Tresses, Slender 66
Lady's Slipper, Large Yellow 126
Lady's Slipper, Showy 101
Lady's Slipper, Small Yellow 126
Lady's Slipper, Stemless or Pink 204
Lady's Thumb 248
Lambkill 220
Larkspur, Dwarf 317
Larkspur, Tall Wild 317
Lathyrus maritimus 314
Laurel, Mountain 212
Laurel, Pale 220

Digitized by Google

PAGE

Laurel, Sheep 220
Lavender, Sea 318
Leek, Wild 46
Leontodon autumnalis 166
Leonurus Cardiaca 290
Lettuce, Wild 146
Liatris scariosa 302
Lilium canadense 134
Lilium philadelphicum 224
Lilium superbum 134
Lily, Cow 132
Lily, Meadow or Field 134
Lily, Turk's Cap 134
Lily, White Water 103
Lily, Wild Orange-red 224
Lily, Wild Yellow 134
Lily, Wood 224
Lily, Yellow Pond 132
Limonium carolinianum 318
Linaria canadensis 316
Linaria vulgaris 160
Linnaea borealis 226
Liparis liliifolia 340
Live-for-ever 254
Liverleaf 270
Liverleaf, Sharp-lobed 270
Liverwort 270
Lizard's Tail 101
Lobelia cardinalis 244
Lobelia, Pale Spiked 290
Lobelia spicata 290
Loosestrife 140
Loosestrife, Four-leaved 140
Loosestrife, Fringed 136
Lousewort 216
Lupine, Wild 274
Lupinus perennis 274
Lychnis alba 50
Lychnis, Evening 50
Lysimachia quadrifolia 140
Lysimachia terrestris 140

Maianthemum canadense 26
Mallow, Common 52

Digitized by Google

PAGE

Mallow, Musk 52
Mallow Rose 265
Mallow, Swamp Rose 265
Malva moschata 52
Malva rotundifolia 52
Mandrake 97
Marigold, Marsh 112
May Apple 97
Mayflower 94
Meadow Beauty 246
Meadow Beauty, Maryland 246
Meadow Rue, Early 338
Meadow Rue, Tall 78
Meadow-sweet 228
Medeola virginiana 120
Melilot, White 86
Melilot, Yellow 181
Melilotus alba 86
Melilotus officinalis 181
Mentha arvensis var. canadensis 298
Mentha piperita 298
Mentha spicata 298
Mercury 34
Mertensia virginica 310
Milfoil 58
Milkweed, Common 236
Milkweed, Four-leaved 242
Milkweed, Poke or Tall 102
Milkweed, Purple 236
Milkweed, Swamp 242
Milkwort, Cross-leaved 264
Milkwort, Field 210
Milkwort, Fringed 210
Mimulus ringens 292
Mint, Wild 298
Mitchella repens 6
Mitella diphylla 10
Mitella nuda 10
Mitrewort 10
Mitrewort, False 10
Mitrewort, Naked 10
Moccasin Flower 204
Monarda didyma 246
Monarda fistulosa 263

Digitized by Google

PAGE

Moneses uniflora 38
Monkey Flower 292
Monotropa Hypopitys 330
Monotropa uniflora 46
Morning-glory, Wild 238
Moss, Flowering 94
Motherwort 290
Mullein, Common 144
Mullein, Moth 144
Mustard, Black 150
Myosotis laxa 313
Myosotis scorpioides 313

Nepeta hederacea 276
New Jersey Tea 100
Nightshade 282
Nymphaea advena 132

Oakesia sessilifolia 122
Oenothera biennis 154
Oenothera fruticosa 154
Oenothera pumila 154
Orchis, Greater Green 64
Orchis, Hooker's 64
Orchis, Large Round-leaved 64
Orchis, Large Purple Fringed 234
Orchis, Prairie White Fringed 104
Orchis, Purple 317
Orchis, Ragged Fringed 62
Orchis, Rein 42
Orchis, Showy 206
Orchis, Smaller Purple Fringed 234
Orchis spectabilis 206
Orchis, White Fringed 104
Orchis, Yellow Fringed 162
Orontium aquaticum 180
Orpine, Garden 254
Oswego Tea 246
Oxalis Acetosella 84
Oxalis corniculata 148
Oxalis violacea 259

Painted Cup 260
Panax quinquefolium 183

Digitized by Google

PAGE
Panax trifolium 20
Parnassia caroliniana 80
Parsnip, Early Meadow 116
Parsnip, Water 102
Parsnip, Wild or Field 116
Partridge Berry 6
Pastinaca sativa 116
Pea, Beach 314
Pea, Partridge 166
Pedicularis canadensis 216
Peltandra virginica 24
Pentstemon hirsutus 259
Pepperbush, Sweet 40
Peppermint 298
Persicaria, Water 248
Phlox divaricata 312
Phlox, Downy 258
Phlox pilosa 258
Phlox subulata 258
Phlox, Wild Blue 312
Phytolacca decandra 92
Pickerel-weed 292
Pigeon Berry 92
Pinesap 330
Pink, Deptford 196
Pink, Grass 222
Pink, Ground or Moss 258
Pink, Indian 190
Pink, Large Marsh 240
Pink, Maiden 196
Pink, Sea 240
Pink, Slender Marsh 240
Pink, Wild 196
Pinxter Flower 208
Pipsissewa 234
Pitcher-plant 202
Plantain, Robin's 274
Plantain, Water 82
Pleurisy Root 183
Podophyllum peltatum 97
Pogonia ophioglossoides 222
Pogona, Rose or Sweet 222
Pokeweed 92
Polygala cruciata 264

Digitized by Google

PAGE

Polygala, Field 210
Polygala, Fringed 210
Polygala paucifolia 210
Polygala sanguinea 210
Polygonatum biflorum 16
Polygonatum commutatum 16
Polygonum amphibium 248
Polygonum pennsylvanicum 248
Polygonum Persicaria 248
Pontederia cordata 292
Potentilla canadensis 114
Prenanthes altissima 332
Primrose, Common Evening 154
Prince's Pine 234
Prunella vulgaris 284
Pussy's Toes 12
Pyrola americana 38
Pyrola elliptica 38
Pyrola, One-flowered 38
Pyrola, Round-leaved 38
Pyxidanthera barbulata 94
Pyxie 94

Queen Anne's Lace 88
Queen of the Prairie 261

Radicula Nasturtium-aquaticum 99
Ragwort, Golden 128
Raspberry, Purple Flowering 230
Rattlesnake Plantain, Downy 62
Rattlesnake-root 332
Rattlesnake-weed 174
Red-root 100
Rheumatism Root 96
Rhexia mariana 246
Rhexia virginica 246
Rhododendron nudiflorum 208
Rhododendron viscosum 100
Rhus Toxicodendron 34
Rhus Vernix 34
Rich-weed 184
Rocket, Yellow 150
Rockrose 182
Rosa blanda 262

Digitized by Google

PAGE

Rosa carolina 262
Rosa humilis 263
Rosa rubiginosa 263
Rosa virginiana 262
Rose, Pasture or Dwarf 262
Rose, Smooth 262
Rose, Swamp 262
Rosemary, Marsh 318
Rubus odoratus 230
Rudbeckia hirta 142
Rudbeckia laciniata 142

Sabatia angularis 240
Sabatia dodecandra 240
Sabatia gracilis 240
Sabatia stellaris 240
Sagittaria latifolia 72
Sambucus canadensis 40
Sambucus racemosa 40
Sanguinaria canadensis 2
Saponaria officinalis 216
Sarracenia purpurea 202
Sarsaparilla, Bristly 36
Sarsaparilla, False or Wild 20
Saururus cernuus 101
Saxifraga virginiensis 12
Saxifrage, Early 12
Scabious, Sweet 56
Scutellaria integrifolia 294
Scutellaria lateriflora 294
Sedum purpureum 254
Self-heal 284
Senecio aureus 128
Senna, Wild 166
Shin Leaf 38
Shooting Star 190
Silene latifolia 50
Silene pennsylvanica 196
Sisyrinchium angustifolium 284
Sium cicutaefolium 102
Skullcap 294
Skullcap, Mad-dog 294
Skunk Cabbage 322
Smartweed 248

Digitized by Google

PAGE

Smilacina racemosa 32
Smilacina stellata 32
Smilax herbacea 130
Snake Mouth 222
Snakeroot, Black 100
Snakeroot, White 68
Sneezeweed 184
Snowberry, Creeping 99
Soapwort 216
Solanum Dulcamara 282
Solomon's Seal 16
Solomon's Seal, False 32
Solomon's Seal, Great 16
Solomon's Seal, Star-flowered 32
Sorrel, Lady's 148
Sorrel, Violet Wood 259
Sorrel, Wood 84
Sparganium americanum 328
Spatter-dock 132
Spearmint 298
Speedwell, Common 314
Spiderwort 313
Spikenard, False 32
Spiraea latifolia 228
Spiraea tomentosa 228
Spiranthes cernua 66
Spiranthes gracilis 66
Spring Beauty 188
Squirrel Corn 4
Star Flower 30
Star Grass 114
Steeple Bush 228
Steironema ciliatum 136
St. John's Wort, Common 158
Streptopus amplexifolius 214
Streptopus roseus 214
Succory 302
Sumach, Poison 34
Sundew, Long-leaved 66
Sundew, Round-leaved 66
Sundrops 154
Sunflower, Swamp 184
Sunflower, Thin-leaved 176
Sweetbrier 263

Digitized by Google

PAGE

Symplocarpus foetidus 322

Tanacetum vulgare 170
Tansy 170
Thalictrum dioicum 338
Thalictrum polygamum 78
Thimbleweed 44
Thoroughwort 68
Tiarella cordifolia 10
Toadflax, Blue 316
Toadflax, Yellow 160
Toothwort 6
Touch-me-not, Pale 156
Touch-me-not, Spotted 156
Tradescantia virginiana 313
Trientalis americana 30
Trifolium agrarium 148
Trifolium arvense 252
Trillium cernuum 28
Trillium erectum 192
Trillium grandiflorum 28
Trillium, Large-flowered 28
Trillium, Nodding 28
Trillium, Painted 28
Trillium, Purple 192
Trillium recurvatum 192
Trillium sessile 192
Trillium, Stemless 192
Trillium undulatum 28
Turtlehead 70
Twayblade, Large 340
Twin-flower 226
Twinleaf 96
Twisted-stalk 214
Twisted-stalk, Clasping-leaved 214
Typha angustifolia 341
Typha latifolia 341

Uvularia grandiflora 122
Uvularia perfoliata 122

Veratrum viride 181
Verbascum Blattaria 144
Verbascum Thapsus 144

Digitized by Google

PAGE

Verbena hastata 296
Vernonia noveboracensis 300
Veronica americana 315
Veronica officinalis 314
Veronica virginica 105
Vervain, Blue 296
Vetch, American 280
Vetch, Cow 280
Vicia americana 280
Vicia Cracca 280
Viola canadensis 98
Viola conspersa 311
Viola lanceolata 98
Viola pallens 97
Viola palmata 310
Viola papilionacea 312
Viola pedata 272
Viola pubescens 180
Viola rotundifolia 179
Viola sagittata 311
Violet, Arrow-leaved 311
Violet, Bird-foot 272
Violet, Canada 98
Violet, Common 312
Violet, Dog 311
Violet, Dog's-tooth 110
Violet, Downy Yellow 180
Violet, Early Blue 310
Violet, Early Yellow 179
Violet, Lance-leaved 98
Violet, Round-leaved 179
Violet, Sweet White 97
Virgin's Bower 60
Wake Robin 98–192
Widow's or Job's Tears 313
Willow-herb, Great 244
Wind-flower 18
Wintergreen 104
Wintergreen, Flowering 210
Wintergreen, Spotted 234

Yarrow 58

Zizia aurea 116

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

This book should be returned to the Library on or before the last date stamped below.

A fine of five cents a day is incurred by retaining it beyond the specified time.

Please return promptly.

DUE JUN 23 '38

JUL 14 1938

DUE AUG 12 '38

DUE JUL 19 '39

DUE MAR -3 '41

DUE AUG 25 '41

DUE NOV -9 '49

DUE NOV 18 '49

DUE MAR -4 '50

Digitized by Google

Digitized by Google